

**Wednesday 5 June 2013 – Afternoon**

**GCSE TWENTY FIRST CENTURY SCIENCE  
ADDITIONAL SCIENCE A**

**A151/02** Modules B4 C4 P4 (Higher Tier)

Candidates answer on the Question Paper.  
A calculator may be used for this paper.

**OCR supplied materials:**  
None

**Other materials required:**

- Pencil
- Ruler (cm/mm)

**Duration:** 1 hour



Candidate forename		Candidate surname	
--------------------	--	-------------------	--

Centre number						Candidate number				
---------------	--	--	--	--	--	------------------	--	--	--	--

**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the boxes above. Please write clearly and in capital letters.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Write your answer to each question in the space provided. Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.

**INFORMATION FOR CANDIDATES**

- Your quality of written communication is assessed in questions marked with a pencil (✎).
- The number of marks is given in brackets [ ] at the end of each question or part question.
- A list of physics equations is printed on page 2.
- The Periodic Table is printed on the back page.
- The total number of marks for this paper is **60**.
- This document consists of **20** pages. Any blank pages are indicated.

## TWENTY FIRST CENTURY SCIENCE EQUATIONS

### Useful relationships

#### The Earth in the Universe

$$\text{distance} = \text{wave speed} \times \text{time}$$

$$\text{wave speed} = \text{frequency} \times \text{wavelength}$$

#### Sustainable energy

$$\text{energy transferred} = \text{power} \times \text{time}$$

$$\text{power} = \text{voltage} \times \text{current}$$

$$\text{efficiency} = \frac{\text{energy usefully transferred}}{\text{total energy supplied}} \times 100\%$$

#### Explaining motion

$$\text{speed} = \frac{\text{distance travelled}}{\text{time taken}}$$

$$\text{acceleration} = \frac{\text{change in velocity}}{\text{time taken}}$$

$$\text{momentum} = \text{mass} \times \text{velocity}$$

$$\text{change of momentum} = \text{resultant force} \times \text{time for which it acts}$$

$$\text{work done by a force} = \text{force} \times \text{distance moved in the direction of the force}$$

$$\text{amount of energy transferred} = \text{work done}$$

$$\text{change in gravitational potential energy} = \text{weight} \times \text{vertical height difference}$$

$$\text{kinetic energy} = \frac{1}{2} \times \text{mass} \times [\text{velocity}]^2$$

#### Electric circuits

$$\text{power} = \text{voltage} \times \text{current}$$

$$\text{resistance} = \frac{\text{voltage}}{\text{current}}$$

$$\frac{\text{voltage across primary coil}}{\text{voltage across secondary coil}} = \frac{\text{number of turns in primary coil}}{\text{number of turns in secondary coil}}$$

#### Radioactive materials


$$\text{energy} = \text{mass} \times [\text{speed of light in a vacuum}]^2$$



- 2 Karen investigates osmosis using potato cylinders. She cuts six potato cylinders, each 50 mm long. She places each potato cylinder in a different concentration of sugar solution. After 60 minutes, she removes the potato cylinders and measures the length of each one.

Concentration of sugar solution in $\text{g}/\text{dm}^3$	% change in length of potato cylinder
0	+6.2
20	+3.0
40	+1.0
60	-1.3
80	-4.4
100	-6.6

- (a) Plot a graph of these results on the axes and draw a straight line of best fit. Four points have been done for you.



[1]

(b) What is the concentration inside the potato cells at the start of the experiment?

Justify your answer.

.....  
.....  
..... [2]

(c) Explain how Karen could improve her experiment to increase confidence in the conclusion.


.....  
.....  
..... [2]

[Total: 5]

**Question 3 begins on page 6**

- 3 Paul studies how plants make glucose by photosynthesis.  
The chemical formula for glucose is  $C_6H_{12}O_6$ .

(a) Write a balanced symbol equation for the process of photosynthesis.



(b) Some of the glucose produced by photosynthesis is used to make other chemicals in plant cells.

One of these other chemicals is the amino acid proline,  $C_5H_9NO_2$ .

From this formula Paul concludes that one molecule of glucose is the only thing needed to make one molecule of proline.

Paul is wrong.

Explain **two** reasons why.


.....

.....

.....

..... [3]

- (c) Paul investigates the rate of photosynthesis at different concentrations of carbon dioxide at three different temperatures. The light intensity is kept constant. The results are plotted on this graph.



The rate of photosynthesis can be affected by several limiting factors.

What is the limiting factor at point **A**?

Explain your answer.

.....

.....

.....

..... [2]

(d) Paul makes these statements about his results.

Two of the statements can be concluded from Paul's results shown in the graph.

The other two statements require further research.

Put a tick (✓) in each row to complete the table.


	<b>Can be concluded</b>	<b>Needs further research</b>
Photosynthesis will slow down if there is not enough carbon dioxide.		
The overall rate of photosynthesis is due to a combination of three factors.		
Plants must take in carbon dioxide by diffusion.		
More than just the carbon dioxide concentration affects the rate of photosynthesis.		

[2]

[Total: 9]



4 Nigella puts a substance into a flame.





She photographs its spectrum.



Nigella thinks that the substance contains sodium compounds and potassium compounds.

Jo thinks it contains sodium compounds, but no potassium compounds.

They look up some spectra in a book.



Who is correct? Explain your answer.

.....

.....

.....

..... [3]

[Total: 3]

5 The halogens are very reactive. This is because of the way that the nucleus holds the electrons around each atom.

(a) The relative atomic mass of one type of chlorine is 37.

This chlorine atom has 17 electrons around the nucleus.

Use this information to decide how many protons and neutrons are in the nucleus of each chlorine atom.

protons ..... neutrons ..... [1]

(b) The electron arrangement is 2.8.7 for a chlorine atom.

The element fluorine is also a halogen.

What is the electron arrangement for a fluorine atom?

..... [1]


(c) When halogens react with metals, the halogen atoms become charged.

Describe how the halogen atoms turn into charged particles.

.....  
.....  
..... [3]

(d) A solution of chlorine reacts with a solution of potassium iodide. A solid is made.

(i) Put state symbols in the boxes for the word equation for this reaction.



[1]

(ii) Write a balanced chemical equation for the reaction.

..... [2]

(iii) Chlorine atoms are very reactive.

The chloride ions in potassium chloride are not very reactive.

Use your understanding of electrons to explain why chloride ions are much less reactive than chlorine atoms.

.....  
 .....  
 ..... [2]

(e) Potassium chloride solution conducts electricity.

Why is this?

Put a tick [✓] in the box next to the best reason.

Potassium is a metal and metals conduct.

Potassium chloride is made of ions.

The solution contains ions which can move.

The water conducts the electricity.

[1]

[Total: 11]

6 In 1869, Mendeleev arranged the known elements into his Periodic Table.

Two new elements were discovered a few years later.

After these discoveries scientists started to agree with Mendeleev’s table.

Suggest why the discovery of these two new elements helped his Periodic Table to become accepted.



*The quality of written communication will be assessed in your answer.*

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

[6]

[Total: 6]

7 Modern cars are built with larger crumple zones than older cars.



Explain why larger crumple zones are more effective than small ones.



*The quality of written communication will be assessed in your answer.*

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

[6]

**[Total: 6]**

- 8 Danny holds a metre ruler above Ellie's open hand. She catches the ruler when he lets go of it without warning. They use the distance that the ruler falls to work out her reaction time.



(a) Here are their results.

Test	Reaction time in seconds
one	0.29
two	0.18
three	0.22
four	0.19
five	0.21

Danny calculates the best estimate of the true value of her reaction time.

He uses  $\frac{0.18 + 0.22 + 0.19 + 0.21}{4} = 0.20\text{s}$ .

Is he correct? Justify your answer.

.....

.....

.....

.....

.....

[2]

- (b) Ellie drinks an energy drink.  
Half an hour later Danny measures Ellie's reaction time again.  
Here are their results.

Test	Reaction time in seconds
six	0.18
seven	0.19
eight	0.16
nine	0.17
ten	0.20

Does drinking the energy drink affect Ellie's reaction time?  
Justify your answer. Use data from the tables.


.....

.....

.....

..... [3]

(c) Danny uses this graph to convert distance fallen by the ruler into a reaction time.



Explain how the graph shows that the ruler is speeding up as it falls.

.....

.....

..... [2]

[Total: 7]







9 A policeman uses a radar gun to measure the speed of a lorry moving towards him.



- (a) He makes two speed measurements of the lorry.  
 The first measurement gives a speed of 31 m/s.  
 The second measurement gives a speed of 25 m/s.  
 The measurements are separated by 1.5 s.  
 Calculate the acceleration of the lorry.

acceleration = ..... m/s<sup>2</sup> [2]


- (b) Which of the distance-time graphs shows the motion of the lorry?  
 The distance is between the lorry and the policeman.




answer ..... [1]

[Total: 3]

10 Jim pushes a heavy crate across a level floor.



(a) Four different forces act on the heavy crate.  
One of them is shown below, acting to the right.



The force from Jim acts to the right.  
Describe the other three forces acting on the crate.

.....

.....

.....

..... [3]

(b) Jim has a weight of 800 N.  
He pushes **on** the crate with a force of 200 N, giving it a steady speed of 0.5 m/s.  
The crate has a weight of 400 N and a mass of 40 kg.  
Draw **one** line to link the direction of the force **on** Jim from the crate to its correct **size**.  
Draw another line to link the **size** to the correct **explanation**.

direction	size	explanation
up	100 N	Force is equal to weight times speed.
left	200 N	The force is half of an interaction pair.
right	400 N	Jim is not leaning his weight against the crate.
down	800 N	The friction equals the weight for steady speed.

[1]

[Total: 4]

END OF QUESTION PAPER

**PLEASE DO NOT WRITE ON THIS PAGE**



**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

# The Periodic Table of the Elements

	1	2	3	4	5	6	7	0		
	7 <b>Li</b> lithium 3	9 <b>Be</b> beryllium 4	11 <b>Na</b> sodium 11	12 <b>C</b> carbon 6	13 <b>Al</b> aluminium 13	14 <b>N</b> nitrogen 7	15 <b>P</b> phosphorus 15	16 <b>O</b> oxygen 8	17 <b>Cl</b> chlorine 17	18 <b>Ar</b> argon 18
	19 <b>K</b> potassium 19	20 <b>Ca</b> calcium 20	23 <b>V</b> vanadium 23	24 <b>Cr</b> chromium 24	25 <b>Mn</b> manganese 25	26 <b>Fe</b> iron 26	27 <b>Co</b> cobalt 27	28 <b>Ni</b> nickel 28	29 <b>Cu</b> copper 29	30 <b>Zn</b> zinc 30
	37 <b>Rb</b> rubidium 37	38 <b>Sr</b> strontium 38	40 <b>Y</b> yttrium 39	41 <b>Zr</b> zirconium 40	42 <b>Nb</b> niobium 41	43 <b>Tc</b> technetium [98]	44 <b>Ru</b> ruthenium 44	45 <b>Rh</b> rhodium 45	46 <b>Pd</b> palladium 46	47 <b>Cd</b> cadmium 48
	55 <b>Cs</b> caesium 55	56 <b>Ba</b> barium 56	57 <b>La*</b> lanthanum 57	72 <b>Hf</b> hafnium 72	73 <b>Ta</b> tantalum 73	74 <b>W</b> tungsten 74	75 <b>Re</b> rhenium 75	76 <b>Os</b> osmium 76	77 <b>Ir</b> iridium 77	78 <b>Pt</b> platinum 78
	87 <b>Fr</b> francium 87	88 <b>Ra</b> radium 88	89 <b>Ac*</b> actinium 89	104 <b>Rf</b> rutherfordium 104	105 <b>Db</b> dubnium 105	106 <b>Sg</b> seaborgium 106	107 <b>Bh</b> bohrium 107	108 <b>Hs</b> hassium 108	109 <b>Mt</b> meitnerium 109	110 <b>Ds</b> darmstadtium 110
	133 <b>Cs</b> caesium 55	137 <b>Ba</b> barium 56	139 <b>La*</b> lanthanum 57	178 <b>Hf</b> hafnium 72	181 <b>Ta</b> tantalum 73	184 <b>W</b> tungsten 74	186 <b>Re</b> rhenium 75	190 <b>Os</b> osmium 76	192 <b>Ir</b> iridium 77	195 <b>Pt</b> platinum 78
	223 <b>Fr</b> francium 87	226 <b>Ra</b> radium 88	227 <b>Ac*</b> actinium 89	261 <b>Rf</b> rutherfordium 104	262 <b>Db</b> dubnium 105	266 <b>Sg</b> seaborgium 106	268 <b>Mt</b> meitnerium 109	271 <b>Ds</b> darmstadtium 110	272 <b>Rg</b> roentgenium 111	201 <b>Hg</b> mercury 80
	131 <b>Xe</b> xenon 54	127 <b>I</b> iodine 53	122 <b>Sb</b> antimony 51	119 <b>Sn</b> tin 50	115 <b>In</b> indium 49	112 <b>Cd</b> cadmium 48	108 <b>Ag</b> silver 47	106 <b>Pd</b> palladium 46	103 <b>Rh</b> rhodium 45	101 <b>Ru</b> ruthenium 44
	[222] <b>Rn</b> radon 86	[210] <b>At</b> astatine 85	[209] <b>Po</b> polonium 84	207 <b>Pb</b> lead 82	204 <b>Tl</b> thallium 81	209 <b>Bi</b> bismuth 83	128 <b>Te</b> tellurium 52	122 <b>Sb</b> antimony 51	79 <b>Se</b> selenium 34	75 <b>As</b> arsenic 33
	4 <b>He</b> helium 2	20 <b>Ne</b> neon 10	40 <b>Ar</b> argon 18	84 <b>Kr</b> krypton 36	131 <b>Xe</b> xenon 54	178 <b>Rn</b> radon 86	19 <b>F</b> fluorine 9	16 <b>O</b> oxygen 8	14 <b>N</b> nitrogen 7	12 <b>C</b> carbon 6
	1 <b>H</b> hydrogen 1	2 <b>He</b> helium 2	3 <b>Li</b> lithium 3	4 <b>Be</b> beryllium 4	5 <b>B</b> boron 5	6 <b>C</b> carbon 6	7 <b>N</b> nitrogen 7	8 <b>O</b> oxygen 8	9 <b>F</b> fluorine 9	10 <b>Ne</b> neon 10

**Key**  
relative atomic mass  
atomic symbol  
name  
atomic (proton) number

Elements with atomic numbers 112-116 have been reported but not fully authenticated

\* The lanthanoids (atomic numbers 58-71) and the actinoids (atomic numbers 90-103) have been omitted.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.