



Tuesday 10 June 2014 – Morning

A2 GCE MATHEMATICS

4733/01 Probability & Statistics 2

QUESTION PAPER

Candidates answer on the Printed Answer Book.

OCR supplied materials:


- Printed Answer Book 4733/01
- List of Formulae (MF1)

Other materials required:

- Scientific or graphical calculator

Duration: 1 hour 30 minutes

MODIFIED LANGUAGE



INSTRUCTIONS TO CANDIDATES

These instructions are the same on the Printed Answer Book and the Question Paper.

- The Question Paper will be found inside the Printed Answer Book.
- Write your name, centre number and candidate number in the spaces provided on the Printed Answer Book. Please write clearly and in capital letters.
- **Write your answer to each question in the space provided in the Printed Answer Book.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.

INFORMATION FOR CANDIDATES

This information is the same on the Printed Answer Book and the Question Paper.

- The number of marks is given in brackets [] at the end of each question or part question on the Question Paper.
- **You are reminded of the need for clear presentation in your answers.**
- The total number of marks for this paper is **72**.
- The Printed Answer Book consists of **12** pages. The Question Paper consists of **4** pages. Any blank pages are indicated.

INSTRUCTION TO EXAMS OFFICER/INVIGILATOR

- Do not send this Question Paper for marking; it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

- 1 The random variable F has the distribution $B(50, 0.7)$. Use a suitable approximation to find $P(F > 40)$. [5]
- 2 The events organiser of a school sends out invitations to 150 people to attend its prize day. From past experience the organiser knows that the number of those who will come to the prize day can be modelled by the distribution $B(150, 0.98)$.

(i) Explain why this distribution cannot be well approximated by either a normal or a Poisson distribution. [3]

(ii) By considering the number of those who do not attend, use a suitable approximation to find the probability that fewer than 146 people attend. [4]

- 3 The random variable G has the distribution $N(\mu, \sigma^2)$. One hundred observations of G are taken. The results are summarised in the following table.

Interval	$G < 40.0$	$40.0 \leq G < 60.0$	$G \geq 60.0$
Frequency	17	58	25

(i) By considering $P(G < 40.0)$, write down an equation involving μ and σ . [2]

(ii) Find a second equation involving μ and σ . Hence calculate values for μ and σ . [4]

(iii) Explain why your answers are only estimates. [1]

- 4 A zoologist investigates the number of snakes found in a given region of land. The zoologist intends to use a Poisson distribution to model the number of snakes.

(i) One condition for a Poisson distribution to be valid is that snakes must occur at constant average rate. State another condition needed for a Poisson distribution to be valid. [1]

Assume now that the number of snakes found in 1 acre of a region can be modelled by the distribution $Po(4)$.

(ii) Find the probability that, in 1 acre of the region, at least 6 snakes are found. [2]

(iii) Find the probability that, in 0.77 acres of the region, the number of snakes found is either 2 or 3. [4]

- 5 A continuous random variable X has probability density function

$$f(x) = \begin{cases} \frac{1}{2}\pi \sin(\pi x) & 0 \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

(i) Show that this is a valid probability density function. [4]

(ii) Sketch the curve $y = f(x)$ and write down the value of $E(X)$. [3]

(iii) Find the value q such that $P(X > q) = 0.75$. [3]

(iv) Write down an expression, including an integral, for $\text{Var}(X)$. (Do not attempt to evaluate the integral.) [2]

(v) A student states that “ X is more likely to occur when x is close to $E(X)$.” Give an improved version of this statement. [1]

- 6 In a city the proportion of inhabitants from ethnic group Z is known to be 0.4. A sample of 12 employees of a large company in this city is obtained and it is found that 2 of the employees are from ethnic group Z . A test is carried out, at the 5% significance level, of whether the proportion of employees from ethnic group Z in this company is less than the proportion in the city as a whole.

(i) State an assumption that must be made about the sample for a significance test to be valid. [1]

(ii) Describe briefly an appropriate way of obtaining the sample. [2]

(iii) Carry out the test. [7]

(iv) A manager believes that the company discriminates against ethnic group Z . Explain whether carrying out the test at the 10% significance level would be more supportive or less supportive of the manager's belief. [2]

- 7 An examination board is developing a new syllabus and wants to know if the question papers are the right length. A random sample of 50 candidates was given a pre-test on a dummy paper. The times, t minutes, taken by these candidates to complete the paper can be summarised by

$$n = 50, \quad \sum t = 4050, \quad \sum t^2 = 329\,800.$$

Assume that times are normally distributed.

(i) Estimate the proportion of candidates that could not complete the paper within 90 minutes. [6]

(ii) Test, at the 10% significance level, whether the mean time for all candidates to complete this paper is 80 minutes. Use a two-tail test. [7]

(iii) Explain whether the assumption that times are normally distributed is necessary in answering

(a) part (i),

(b) part (ii). [2]

Question 8 begins on page 4.

- 8 The random variable W has the distribution $\text{Po}(\lambda)$. A significance test is carried out of the null hypothesis $H_0: \lambda = 3.60$, against the alternative hypothesis $H_1: \lambda < 3.60$. The test is based on a single observation of W . The critical region is $W = 0$.

(i) Find the significance level of the test. [2]

(ii) When $\lambda = \lambda_0$, the probability that the test results in a Type II error is 0.8. Find the value of λ_0 . [4]

END OF QUESTION PAPER



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.