

GCE

Mathematics (MEI)

Unit 4757: Further Applications of Advanced Mathematics

Advanced GCE

Mark Scheme for June 2018

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2018

Annotations and abbreviations

Annotation in scoris	Meaning
✓and ×	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
<u>۸</u>	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark	Meaning
scheme	
scheme E1	Mark for explaining
scheme E1 G1	Mark for explaining Mark for a correct feature on a graph
scheme E1 G1 M1 dep*	Mark for explaining Mark for a correct feature on a graph Method mark dependent on a previous mark, indicated by *
scheme E1 G1 M1 dep* cao	Mark for explaining Mark for a correct feature on a graph Method mark dependent on a previous mark, indicated by * Correct answer only
scheme E1 G1 M1 dep* cao oe	Mark for explaining Mark for a correct feature on a graph Method mark dependent on a previous mark, indicated by * Correct answer only Or equivalent
scheme E1 G1 M1 dep* cao oe rot	Mark for explaining Mark for a correct feature on a graph Method mark dependent on a previous mark, indicated by * Correct answer only Or equivalent Rounded or truncated
scheme E1 G1 M1 dep* cao oe rot soi	Mark for explaining Mark for a correct feature on a graph Method mark dependent on a previous mark, indicated by * Correct answer only Or equivalent Rounded or truncated Seen or implied
scheme E1 G1 M1 dep* cao oe rot soi www	Mark for explaining Mark for a correct feature on a graph Method mark dependent on a previous mark, indicated by * Correct answer only Or equivalent Rounded or truncated Seen or implied Without wrong working
scheme E1 G1 M1 dep* cao oe rot soi www	Mark for explaining Mark for a correct feature on a graph Method mark dependent on a previous mark, indicated by * Correct answer only Or equivalent Rounded or truncated Seen or implied Without wrong working

Subject-specific Marking Instructions for GCE Mathematics Pure strand

a Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded

b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks according to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) you should contact your Team Leader.

c The following types of marks are available.

Μ

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, eg by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

Α

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

В

Mark for a correct result or statement independent of Method marks.

Ε

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

Mark Scheme

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep *' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise. Candidates are expected to give numerical answers to an appropriate degree of accuracy, with 3 significant figures often being the norm. Small variations in the degree of accuracy to which an answer is given (e.g. 2 or 4 significant figures where 3 is expected) should not normally be penalised, while answers which are grossly over- or under-specified should normally result in the loss of a mark. The situation regarding any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. If in doubt, contact your Team Leader.
- g Rules for replaced work

If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

h For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A mark in the question.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

Mark Scheme

1	(i)	Find a common point.		
		e.g. set $z = 0$ then solve $x + 2y = 11$, $2x - y = 7$ simultaneously.	M1	Find a common point
		\Rightarrow (5, 3, 0)	AI	
		Find the vector product	M1	Evaluation of vector product
		$\Rightarrow n_1 = \begin{pmatrix} 1 \\ 2 \\ -2 \end{pmatrix}, n_2 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$ $\Rightarrow n_1 \times n_2 = \begin{pmatrix} 0 \\ -5 \\ -5 \end{pmatrix} \equiv \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$ $\Rightarrow \frac{x-5}{x-5} = \frac{y-3}{x-5} = \frac{z}{x-5} \text{ of } e.g. x = 5, y = z+3$	A1 A1	Or finding a second point and using it to find direction of <i>L</i>
			[5]	
	(ii)	2a-1+1-7 $ 2a-7 $	M1	Using formula for distance (one sufficient;
		$d_1 = \left \frac{2u + 1 + 1}{\sqrt{1^2 + 2^2 + 1^2}} \right = \left \frac{2u + 1}{\sqrt{6}} \right $		<i>modulus not required;</i> <i>M0 if neither 11 nor 7 used)</i>
		$d_{2} = \left \frac{a + 2 - 2 - 11}{\sqrt{1^{2} + 2^{2} + 2^{2}}} \right = \left \frac{a - 11}{\sqrt{9}} \right = \left \frac{a - 11}{3} \right $	A1	Both distances correct (modulus not required)
		$\Rightarrow \frac{2a-7}{\sqrt{6}} = \frac{a-11}{3} \Rightarrow a = \frac{21-11\sqrt{6}}{6-\sqrt{6}} = 2-\frac{3}{2}\sqrt{6}$	M1	Finding two equations (or squaring)
		$\mathbf{Or} \ \frac{2a-7}{\sqrt{6}} = \frac{11-a}{3} \Longrightarrow a = 2 + \frac{3}{2}\sqrt{6}$	A1	Accept (art) -1.67, 5.67 or $\frac{8 \pm \sqrt{216}}{4}$ etc
			[4]	

(iii)	$1+4-14 \neq 11 \text{ but } 2-2+7=7$ $1+0+10=11 \text{ but } 2-0-5 \neq 7$ BC intersects Q at B only; and P at C only. B and C are not on L so BC does not intersect L BC is not parallel to Q (or P), so BC is not parallel to L	B1 B1	www Needs evidence of substitution Convincingly explain why BC does not intersect <i>L</i> and is not parallel to <i>L</i>
(iv)	$\mathbf{d} = \mathbf{C}\mathbf{B} = \begin{pmatrix} 1\\2\\7 \end{pmatrix} - \begin{pmatrix} 1\\0\\-5 \end{pmatrix} = \begin{pmatrix} 0\\2\\12 \end{pmatrix}, \mathbf{e} = \begin{pmatrix} 0\\1\\1 \end{pmatrix}$ $\mathbf{d} \times \mathbf{e} = \begin{pmatrix} 0\\2\\12 \end{pmatrix} \times \begin{pmatrix} 0\\1\\1 \end{pmatrix} = \begin{pmatrix} -10\\0\\0 \end{pmatrix}, \mathbf{d} \times \mathbf{e} = 10$ $\begin{pmatrix} 4\\0 \end{pmatrix}$	[2] B1 M1 A1 ft	For $\begin{pmatrix} 0 \\ 2 \\ 12 \end{pmatrix}$ Vector product of directions Evaluation of vector product
	$\mathbf{a} \cdot \mathbf{b} = \begin{bmatrix} 1 \\ -7 \end{bmatrix} \Rightarrow (\mathbf{a} \cdot \mathbf{b}).(\mathbf{d} \times \mathbf{e})$ $\left \frac{(\mathbf{a} \cdot \mathbf{b}).(\mathbf{d} \times \mathbf{e})}{ \mathbf{d} \times \mathbf{e} } \right = 4$	M1 A1	Using correct formula to evaluate the distance cao

Mark Scheme

June 2018

(v)	$\mathbf{n} = \begin{pmatrix} 2 \\ -1 \end{pmatrix} \Rightarrow CE$ has equation	(CE is perpendicular to	M1	
	$\mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ -5 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1+2\lambda \\ -\lambda \\ -5+\lambda \end{pmatrix}$	(Equation of CE)	A1 A1	Alternative scheme M1 Finding distance from C to Q A1 Distance is $\frac{5}{3}\sqrt{6}$ A1 Length of CE is $\frac{10}{3}\sqrt{6}$ (2)
	Meets Q when $2+4\lambda + \lambda - 5 + \lambda = 7$ $\Rightarrow 6\lambda = 10 \Rightarrow \lambda = \frac{5}{3}$ E corresponds to $\lambda = \frac{10}{3}$	(Equation for λ)		A1 CE = $(\pm)\frac{10}{3}\begin{bmatrix} -1\\1 \end{bmatrix}$ or E correct
	So E has coordinates $\left(\frac{23}{3}, -\frac{10}{3}, -\frac{5}{3}\right)$		A1 M1	Appropriate scalar triple product stated
	Volume of tetrahedron $=\frac{1}{6} OE.(OB \times OC) $			(any 3 non-coplanar edges) (1/6 not required)
	$ \left \begin{array}{c} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 $		A1 ft	Correct vectors substituted into correct formula (including 1/6)
	$ \begin{vmatrix} -\frac{1}{6} \\ -\frac{1}{6} \\ -\frac{5}{3} \end{vmatrix} \cdot \begin{vmatrix} 2 \\ 7 \end{vmatrix} \times \begin{vmatrix} 0 \\ -5 \\ \end{vmatrix} = \begin{vmatrix} -\frac{1}{6} \\ -\frac{5}{3} \\ -\frac{5}{3} \end{vmatrix} \cdot \begin{vmatrix} 12 \\ -2 \\ -2 \end{vmatrix} $		A1 ft	Evaluation of vector product cao Accept 18.9
	$=\frac{1}{6}\left \left(-\frac{230}{3}-\frac{120}{3}+\frac{10}{3}\right)\right =\frac{340}{18}$			
	\Rightarrow Volume of solid = $\frac{170}{9}$		_	
			[8]	

2	(i)	$\frac{\partial z}{\partial x} = 1 + 8xy, \frac{\partial z}{\partial y} = 4x^2 - 4y$	B1B1	
		At P (1, 1, 5), $\frac{\partial z}{\partial x} = 9$, $\frac{\partial z}{\partial y} = 0$	B1	
		Tangent plane at is $z - c = \frac{\partial z}{\partial x} \Big _{x \to y} (x - a) + \frac{\partial z}{\partial y} \Big _{x \to y} (y - b)$	M1	or $z=9x+d$ or $9x-z=d$
		$\Rightarrow z - 5 = 9(x - 1) + 0(x - 1) \Rightarrow z - 5 = 9(x - 1) \Rightarrow z = 9x - 4$	A1	AG
			[5]	
	(ii)	For tangent plane to be $z = 9x + k$, $\frac{\partial z}{\partial x} = 9$ and $\frac{\partial z}{\partial y} = 0$	M1	
		$\frac{\partial z}{\partial y} = 4x^2 - 4y = 0 \Longrightarrow y = x^2,$		
		$\frac{\partial z}{\partial x} = 1 + 8xy \Longrightarrow 1 + 8x^3 = 9 \Longrightarrow x = 1$	A1	or $y=1$
		$\Rightarrow y = 1 \Rightarrow z = 5$ So $z = 9x + k$ only at (1, 1, 5) $\Rightarrow k = -4$	A1	AG Must mention −4
			[3]	
	(iii)	$\delta z \approx \frac{\partial z}{\partial x} \Big _{p} \delta x + \frac{\partial z}{\partial y} \Big _{p} \delta y$ $\Rightarrow b = 9a + 0 \Rightarrow b = 9a$ ($\delta z \approx 9\delta x + 0\delta y$)	M1 A1 ft A1	Alternative schemeM1 Substituting $x = y = 1 + a, z = 5 + b$ A1 For $b = 9a + 10a^2 + 4a^3$ A1 For $b \approx 9a$ since a is small
			ျပ	

(iv)	$\frac{\partial z}{\partial x} = 1 \text{ and } \frac{\partial z}{\partial y} = 16$ $\Rightarrow 1 + 8xy = 1 \Rightarrow xy = 0 \Rightarrow x = 0 \text{ or } y = 0$	M1	Allow M1 for -1 , -16 M0 for λ , 16λ (unless $\lambda = \pm 1$ seen later)
	$\Rightarrow 4x^{2} - 4y = 16 \Rightarrow x^{2} = y + 4$ $x = 0 \Rightarrow y = -4, z = -30$ i.e. $(0, -4, -30)$ $y = 0 \Rightarrow x = \pm 2, z = 0, 4$ i.e. $(2, 0, 4)$ and $(-2, 0, 0)$	A1 A1A1	SC If M1A0 then SC A1 for all <i>x</i> and <i>y</i> values correct
(V)	$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial y} = 0 \Rightarrow 1 + 8xy = 0, 4x^2 - 4y = 0$ $\Rightarrow y = x^2, 1 + 8xy = 0 \Rightarrow x^3 = -\frac{1}{8} \Rightarrow x = -\frac{1}{2}$ $\Rightarrow y = \frac{1}{4}, z = -\frac{1}{2} + \frac{1}{4} - \frac{1}{8} + 2 = \frac{13}{8}$ $x = -\frac{1}{2} \text{ is the only root of the equation } 1 + 8x^3 = 0$ So only one stationary point. Set $x = -\frac{1}{2} \Rightarrow z = \frac{3}{2} + y - 2y^2$ Set $y = \frac{1}{4} \Rightarrow z = x + x^2 + \frac{15}{8}$ One section has a maximum and the other has a minimum meaning that the stationary point is a saddle point	[4] M1 M1 A1 A1 A1 A1 A1 A1 A1 A1 A1 [9]	For $\frac{\partial z}{\partial x} = \frac{\partial z}{\partial y} = 0$ Obtaining an equation in <i>x</i> (or y) only AG For $x = -\frac{1}{2}$ (or $y = \frac{1}{4}$) AG For other two coordinates (some evidence required for <i>z</i>) AG Must refer to this being the only solution (or the only stationary point) Finding cross-sections (one sufficient) (M0 for $x = 0$ and $y = 0$ used) Correct diagrams sufficient

3	(a)		Formula for surface area = $\int_{x_1}^{x_2} 2\pi y ds$	M1	Used with equation of a circle
			$ds = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \qquad or ds = \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta$	M 1	Used
			$x^{2} + y^{2} = a^{2} \Rightarrow y = \sqrt{a^{2} - x^{2}}, or x = a\cos\theta, y = a\sin\theta$ $\frac{dy}{dx} = -\frac{x}{y} \qquad \qquad or \frac{dx}{d\theta} = -a\sin\theta, \frac{dy}{d\theta} = a\cos\theta$	B1	(in any form)
			$S = 2\int_{0}^{a} 2\pi y \sqrt{1 + \frac{x^2}{y^2}} dx \qquad or S = \int_{0}^{\pi} 2\pi a \sin \theta \sqrt{a^2 \sin^2 \theta + a^2 \cos^2 \theta} d\theta$	A1	Correct integral expression for <i>S</i> including limits
			$= 4\pi a \int_{0}^{a} dx \qquad = 2\pi a^{2} \int_{0}^{\pi} \sin\theta d\theta$	A1 A1	For $y ds = a dx$ or $y ds = a^2 \sin \theta d\theta$
			$=4\pi a^2$		AG
				[6]	
	(b)	(i)	$y = 0 \Longrightarrow 3t^3 = 4t$	M1	
			$t \neq 0 \Longrightarrow t = \pm \frac{2}{3}\sqrt{3}$	A1	Accept $\pm \frac{2}{\sqrt{3}}$ (A0 if $t = 0$ included)
				[2]	
		(ii)	$x = 6t^2 \Longrightarrow \dot{x} = 12t$	B1	
			$y = 4t - 3t^3 \Longrightarrow \dot{y} = 4 - 9t^2$	B1 M1	\mathbf{O} the stitution in term $\mathbf{i}^2 + \mathbf{i}^2$
			$\dot{x}^2 + \dot{y}^2 = \left(4 + 9t^2\right)^2$	A1	Substituting into $x + y$
			$\Rightarrow s = \int_{0}^{\frac{2}{3}\sqrt{3}} (4+9t^{2}) dt = \left[4t+3t^{3}\right]_{0}^{\frac{2}{3}\sqrt{3}} = \frac{2}{3}\sqrt{3}(4+4) = \frac{16}{3}\sqrt{3}$	M1 A1	For $\int \sqrt{\dot{x}^2 + \dot{y}^2} dt$ (limits not required)
				[6]	

June 2018

	(iiii)		B1	
	(,	$\ddot{r} - 12$ $\ddot{v}18t(6)$		
		x = 12, y = 10i(-0)	М1	Lising formula for o or K
		$(\dot{x}^2 + \dot{x}^2)^{3/2}$ $(A + 0A^2)^3$ $(A + 0A^2)^3$		Correct substitution (algebraic or numerical)
		$a = \frac{(x + y)}{(x + y)} = \frac{(4 + 9i)}{(4 + 9i)} = \frac{(4 + 9i)}{(4 + 9i)}$	AI	Correct substitution (algebraic of numerical)
		$p = \frac{1}{10000000000000000000000000000000000$		
		$\frac{1}{12}\left(1-\frac{1}{2}\right) = 100$		
		$=\frac{\left(4+9t^{2}\right)^{3}}{-12\left(4+9t^{2}\right)}=-\frac{1}{12}\left(4+9t^{2}\right)^{2}$		
			A1	
		1 25		llsed
		When $t = \frac{1}{2} \rho = (-)\frac{1}{12}$	М1	0300
		3 12		
			D4	
		Centre is $\mathbf{c} = \mathbf{r} + \rho \hat{\mathbf{n}}$	BI	
		When $t = \frac{1}{3}$, $\mathbf{r} = \begin{pmatrix} \frac{2}{3} \\ \frac{11}{9} \end{pmatrix}$	M1	Finding $\frac{dy}{dx}$
		$\tan w = -\frac{4-9t^2}{3} - \frac{3}{3}$		Correct unit normal
		$\tan \psi = \frac{1}{12t} \frac{1}{4}$	A 1	
		$\Rightarrow \hat{\mathbf{n}} = (\pm) \begin{pmatrix} -\frac{3}{5} \\ \frac{4}{5} \end{pmatrix}$	A1 A1	
		$\Rightarrow \mathbf{c} = \begin{pmatrix} \frac{2}{3} \\ \frac{11}{9} \end{pmatrix} - \frac{25}{12} \begin{pmatrix} -\frac{3}{5} \\ \frac{4}{5} \end{pmatrix} = \begin{pmatrix} \frac{23}{12} \\ -\frac{4}{9} \end{pmatrix}$		
			[10]	

4	(i) (ii)	(A) (B)	Order 2: 4,11, 14 Order 4: 2, 7, 8, 13 1 2 4 7 8 11 13 14 1 4 2 4 4 2 4 2 1, 2, 4, 8 Order 2: {1, 4}, {1, 11}, {1, 14} Order 4: {1, 2, 4, 8}, {1, 4, 7, 13}	B4 [4] B2 [2] B4	B3 one error, B2 2 errors, B1 3 errors B1 for one omission B1 one correct of order 2 B1 one correct of order 4 B1 any 4 correct
					B1 all 5 correct with no extras Ignore {1} if included
	(!!!)			[4]	
	(111)		 (A) Yes. Closed 0 is the identity element Each element has an inverse 1,7 2,6 and 3,5 are inverse pairs, 4 is self inverse Not isomorphic e.g. G has 4 self inverse elements, this has 2 (B) No e.g. 3 × 3 = 0 so set not closed under this operation 	B1 B1 B1 B1 B1 B1 B1	Giving inverses correctly (0 not needed) With reason Reason

Mark Scheme

June 2018

4757	Mark Scheme		June 2018
	(C) Yes $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ is the identity All have inverses $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ is the inverse of $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$	B1 B1 B1 B1	(or vice-versa) With reason
	All other elements are self inverse. Not isomorphic as this one has 6 self-inverse elements	B1 B1	
		[14]	

5	(i)	Lights always change – they are never on for consecutive minutes. The run length is therefore 0 for each	B1 B1 [2]	Accept run length 0 or 1
	(ii)	Finding powers of <i>P</i> ₁ 0, 0.27, 0.162, 0.2091	M1 A1 A1A1 [4]	If M1A0 then SC A1 for three of 0, 0.27, 0.162, 0.2091 regardless of times given
	(iii)	0.162 < <i>a</i> < 0.2091	M1 A1 ft	For lower limit 0 or 0.162 and upper limit 0.27 or 0.2091
	(iv)	$\mathbf{P}_{1}^{4} = \begin{pmatrix} 0.2091 & 0.1976 & 0.201 & 0.1762 \\ 0.329 & 0.3332 & 0.257 & 0.2492 \\ 0.2088 & 0.2054 & 0.2529 & 0.256 \\ 0.2531 & 0.2638 & 0.2891 & 0.3186 \end{pmatrix}$ probabilities for $t = 1$ are 0, 0.2, 0.3, 0.5 P (same at $t = 1$ and 5) = $0.2 \times 0.3332 + 0.3 \times 0.2529 + 0.5 \times 0.3186 = 0.302$ \Rightarrow P (different) = 0.698	[2] M1 A1 M1 M1 A1 A1	Using diagonal elements from P ⁴ . Diagonal elements correct Using probabilities from 2 nd minute Method for P(same) or P(different)
	(v)	$P_{2} = \begin{pmatrix} A & B & C & D \\ A & (0 & 0.5 & 0 & 0) \\ B & (1 & 0 & 0.5 & 0) \\ C & (1 & 0 & 0.5 & 0) \\ 0 & 0.5 & 0 & 1 \\ D & (0 & 0 & 0.5 & 0) \end{pmatrix}$	B2	B1 one error
	(vi)	(A) There are no absorbing states.(B) A and D are reflecting barriers	121 B1 B1B1	Maximum B1 if any extras
			[¹]	

Mark Scheme

(vii)	Algebraically: $\frac{1}{6}, \frac{1}{3}, \frac{1}{3}, \frac{1}{6}$	<i>Alternative scheme</i> M1 Finding large even and large odd powers M1 Averaging over even and odd powers A4 Answers, which must be given as fractions	M1 M1 A4	Using column vector Creating and solving equations (needs to use $a+b+c+d=1$) A1 for each
			[6]	

OCR (Oxford Cambridge and RSA Examinations) The Triangle Building Shaftesbury Road Cambridge CB2 8EA

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998 Facsimile: 01223 552627 Email: <u>general.qualifications@ocr.org.uk</u>

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee Registered in England Registered Office; The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA Registered Company Number: 3484466 OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations) Head office Telephone: 01223 552552 Facsimile: 01223 552553 Cambridge



© OCR 2018