

**ADVANCED GCE  
MATHEMATICS**

**4726/01**

Further Pure Mathematics 2

**FRIDAY 23 MAY 2008**

Morning  
Time: 1 hour 30 minutes

**Additional materials:** Answer Booklet (8 pages)  
List of Formulae (MF1)

**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.


**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is 72.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of 4 printed pages.

- 1 It is given that  $f(x) = \frac{2ax}{(x-2a)(x^2+a^2)}$ , where  $a$  is a non-zero constant. Express  $f(x)$  in partial fractions. [5]

2



The diagram shows the curve  $y = f(x)$ . The curve has a maximum point at  $(0, 5)$  and crosses the  $x$ -axis at  $(-2, 0)$ ,  $(3, 0)$  and  $(4, 0)$ . Sketch the curve  $y^2 = f(x)$ , showing clearly the coordinates of any turning points and of any points where this curve crosses the axes. [5]

- 3 By using the substitution  $t = \tan \frac{1}{2}x$ , find the exact value of

$$\int_0^{\frac{1}{2}\pi} \frac{1}{2 - \cos x} dx,$$

giving the answer in terms of  $\pi$ . [6]

- 4 (i) Sketch, on the same diagram, the curves with equations  $y = \operatorname{sech} x$  and  $y = x^2$ . [3]
- (ii) By using the definition of  $\operatorname{sech} x$  in terms of  $e^x$  and  $e^{-x}$ , show that the  $x$ -coordinates of the points at which these curves meet are solutions of the equation

$$x^2 = \frac{2e^x}{e^{2x} + 1}. \quad [3]$$

- (iii) The iteration

$$x_{n+1} = \sqrt{\frac{2e^{x_n}}{e^{2x_n} + 1}}$$

can be used to find the positive root of the equation in part (ii). With initial value  $x_1 = 1$ , the approximations  $x_2 = 0.8050$ ,  $x_3 = 0.8633$ ,  $x_4 = 0.8463$  and  $x_5 = 0.8513$  are obtained, correct to 4 decimal places. State with a reason whether, in this case, the iteration produces a 'staircase' or a 'cobweb' diagram. [2]

- 5 It is given that, for  $n \geq 0$ ,

$$I_n = \int_0^{\frac{1}{4}\pi} \tan^n x dx.$$

- (i) By considering  $I_n + I_{n-2}$ , or otherwise, show that, for  $n \geq 2$ ,

$$(n-1)(I_n + I_{n-2}) = 1. \quad [4]$$

- (ii) Find  $I_4$  in terms of  $\pi$ . [4]

6 It is given that  $f(x) = 1 - \frac{7}{x^2}$ .

(i) Use the Newton-Raphson method, with a first approximation  $x_1 = 2.5$ , to find the next approximations  $x_2$  and  $x_3$  to a root of  $f(x) = 0$ . Give the answers correct to 6 decimal places. [3]

(ii) The root of  $f(x) = 0$  for which  $x_1, x_2$  and  $x_3$  are approximations is denoted by  $\alpha$ . Write down the exact value of  $\alpha$ . [1]

(iii) The error  $e_n$  is defined by  $e_n = \alpha - x_n$ . Find  $e_1, e_2$  and  $e_3$ , giving your answers correct to 5 decimal places. Verify that  $e_3 \approx \frac{e_2^3}{e_1^2}$ . [3]

7 It is given that  $f(x) = \tanh^{-1}\left(\frac{1-x}{2+x}\right)$ , for  $x > -\frac{1}{2}$ .


(i) Show that  $f'(x) = -\frac{1}{1+2x}$ , and find  $f''(x)$ . [6]

(ii) Show that the first three terms of the Maclaurin series for  $f(x)$  can be written as  $\ln a + bx + cx^2$ , for constants  $a, b$  and  $c$  to be found. [4]

8 The equation of a curve, in polar coordinates, is

$$r = 1 - \sin 2\theta, \quad \text{for } 0 \leq \theta < 2\pi.$$

(i)



The diagram shows the part of the curve for which  $0 \leq \theta \leq \alpha$ , where  $\theta = \alpha$  is the equation of the tangent to the curve at  $O$ . Find  $\alpha$  in terms of  $\pi$ . [2]

(ii) (a) If  $f(\theta) = 1 - \sin 2\theta$ , show that  $f\left(\frac{1}{2}(2k+1)\pi - \theta\right) = f(\theta)$  for all  $\theta$ , where  $k$  is an integer. [3]

(b) Hence state the equations of the lines of symmetry of the curve

$$r = 1 - \sin 2\theta, \quad \text{for } 0 \leq \theta < 2\pi. \quad [2]$$


(iii) Sketch the curve with equation

$$r = 1 - \sin 2\theta, \quad \text{for } 0 \leq \theta < 2\pi.$$

State the maximum value of  $r$  and the corresponding values of  $\theta$ . [4]

- 9 (i) Prove that  $\int_0^N \ln(1+x) dx = (N+1) \ln(N+1) - N$ , where  $N$  is a positive constant. [4]

(ii)



The diagram shows the curve  $y = \ln(1+x)$ , for  $0 \leq x \leq 70$ , together with a set of rectangles of unit width.

- (a) By considering the areas of these rectangles, explain why

$$\ln 2 + \ln 3 + \ln 4 + \dots + \ln 70 < \int_0^{70} \ln(1+x) dx. \quad [2]$$

- (b) By considering the areas of another set of rectangles, show that

$$\ln 2 + \ln 3 + \ln 4 + \dots + \ln 70 > \int_0^{69} \ln(1+x) dx. \quad [3]$$

- (c) Hence find bounds between which  $\ln(70!)$  lies. Give the answers correct to 1 decimal place. [3]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.