

**ADVANCED GCE
MATHEMATICS (MEI)**

Differential Equations

THURSDAY 12 JUNE 2008

4758/01

Morning

Time: 1 hour 30 minutes

Additional materials (enclosed): None

Additional materials (required):

Answer Booklet (8 pages)

Graph paper

MEI Examination Formulae and Tables (MF2)

INSTRUCTIONS TO CANDIDATES

- Write your name in capital letters, your Centre Number and Candidate Number in the spaces provided on the Answer Booklet.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Answer any **three** questions.
- You are permitted to use a graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is **72**.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.

This document consists of **4** printed pages.

- 1 Fig. 1 shows a particle of mass 2 kg suspended from a light vertical spring. At time t seconds its displacement is x m below its equilibrium level and its velocity is v m s⁻¹ vertically downwards. The forces on the particle are

- its weight, $2g$ N
- the tension in the spring, $8(x + 0.25g)$ N
- the resistance to motion, $2kv$ N where k is a positive constant.




Fig. 1

- (i) Use Newton's second law to write down the equation of motion for the particle, justifying the signs of the terms. Hence show that the displacement is described by the differential equation

$$\frac{d^2x}{dt^2} + k\frac{dx}{dt} + 4x = 0. \quad [4]$$

The particle is initially at rest with $x = 0.1$.

- (ii) In the case $k = 0$, state the general solution of the differential equation. Find the solution, subject to the given initial conditions. [4]
- (iii) In the case $k = 2$, find the solution of the differential equation, subject to the given initial conditions. Sketch a graph of the solution for $t \geq 0$. [11]
- (iv) Find the range of values of k for which the system is over-damped. Sketch a possible graph of the solution in such a case. [5]
- 2 The radioactive substance X decays into the substance Y, which in turn decays into Z. At time t hours the masses, in grams, of X, Y and Z are denoted by x , y and z respectively.

Initially there is 8 g of X and there is no Y or Z present.

The differential equation modelling the decay of X is $\frac{dx}{dt} = -2x$.

- (i) Find x in terms of t . [3]

The differential equation modelling the amount of Y is $\frac{dy}{dt} = 2x - y$.

- (ii) Using your expression for x found in part (i), solve this equation to find y in terms of t . [9]

- (iii) Show that $y > 0$ for $t > 0$. Sketch a graph of y for $t \geq 0$. [5]

The differential equation modelling the amount of Z is $\frac{dz}{dt} = y$.

- (iv) Without solving this equation, show that $x + y + z = 8$. Hence show that $z = 8(1 - e^{-t})^2$. [5]

- (v) Calculate the time required for 99% of the total mass to become substance Z. [2]

- 3 The differential equation $t \frac{dy}{dt} + ky = t$, where k is a constant, is to be solved for $t \geq 1$, subject to the condition $y = 0$ when $t = 1$.

(i) When $k \neq -1$, find the solution for y in terms of t and k . [10]

(ii) Sketch a graph of the solution for $k = 2$. [2]

(iii) When $k = -1$, find the solution for y in terms of t . [5]

Now consider the differential equation $t \frac{dy}{dt} - \sin y = t$, subject to the condition $y = 0$ when $t = 1$. This is to be solved by Euler's method. The algorithm is given by $t_{r+1} = t_r + h$, $y_{r+1} = y_r + h\dot{y}_r$.

(iv) Using a step length of 0.1, perform two iterations of the algorithm to estimate the value of y when $t = 1.2$. [4]

If the algorithm is carried out with a step length of 0.05, the estimate for y when $t = 1.2$ is $y \approx 0.2138$.

(v) Explain with a reason which of these two estimates for y when $t = 1.2$ is likely to be more accurate. Hence, or otherwise, explain whether these estimates are likely to be overestimates or underestimates. [3]

- 4 The simultaneous differential equations

$$\begin{aligned} \frac{dx}{dt} &= 4x - 6y - 9 \sin t, \\ \frac{dy}{dt} &= 3x - 5y - 7 \sin t, \end{aligned}$$

are to be solved.

(i) Show that $\frac{d^2x}{dt^2} + \frac{dx}{dt} - 2x = -9 \cos t - 3 \sin t$. [6]

(ii) Find the general solution for x . [9]

(iii) Hence find the corresponding general solution for y . [3]

It is given that x is bounded as $t \rightarrow \infty$.

(iv) Show that y is also bounded as $t \rightarrow \infty$. [2]

(v) Given also that $y = 0$ when $t = 0$, find the particular solutions for x and y . Write down the expressions for x and y as $t \rightarrow \infty$. [4]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.