

**ADVANCED GCE
MATHEMATICS**

4730/01

Mechanics 3

THURSDAY 17 JANUARY 2008

Afternoon

Time: 1 hour 30 minutes

Additional materials: Answer Booklet (8 pages)
List of Formulae (MF1)

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of 4 printed pages.


- 1 A smooth horizontal surface lies in the x - y plane. A particle P of mass 0.5 kg is moving on the surface with speed 5 m s^{-1} in the x -direction when it is struck by a horizontal blow whose impulse has components -3.5 N s and 2.4 N s in the x -direction and y -direction respectively.

- (i) Find the components in the x -direction and the y -direction of the velocity of P immediately after the blow. Hence show that the speed of P immediately after the blow is 5.2 m s^{-1} . [4]

P is struck by a second horizontal blow whose impulse is \mathbf{I} .

- (ii) Given that P 's direction of motion immediately after this blow is parallel to the x -axis, write down the component of \mathbf{I} in the y -direction. [2]


2



Two uniform rods AB and BC , each of length 2 m , are freely jointed at B . The weights of the rods are $W \text{ N}$ and 50 N respectively. The end A of AB is hinged at a fixed point. The rods AB and BC make angles $\tan^{-1}(\frac{3}{4})$ and β respectively with the downward vertical, and are held in equilibrium in a vertical plane by a horizontal force of magnitude 75 N acting at C (see diagram).

- (i) By taking moments about B for BC , show that $\tan \beta = 3$. [3]
- (ii) Write down the horizontal and vertical components of the force acting on AB at B . [2]
- (iii) Find the value of W . [4]

3




Two uniform smooth spheres A and B , of equal radius, have masses 6 kg and 3 kg respectively. They are moving on a horizontal surface when they collide. Immediately before the collision the velocity of A has components 4 m s^{-1} along the line of centres towards B , and $v\text{ m s}^{-1}$ perpendicular to the line of centres. B is moving with speed 8 m s^{-1} along the line of centres towards A (see diagram). The coefficient of restitution between the spheres is e .

- (i) Find, in terms of e , the component of the velocity of A along the line of centres immediately after the collision. [5]
- (ii) Given that the speeds of A and B are the same immediately after the collision, and that $3e^2 = 1$, find v . [4]
- 4 A particle of mass $m\text{ kg}$ is released from rest at a fixed point O and falls vertically. The particle is subject to an upward resisting force of magnitude $0.49mv\text{ N}$ where $v\text{ m s}^{-1}$ is the velocity of the particle when it has fallen a distance of $x\text{ m}$ from O .
- (i) Write down a differential equation for the motion of the particle, and show that the equation can be written as $\left(\frac{20}{20-v} - 1\right)\frac{dv}{dx} = 0.49$. [5]
- (ii) Hence find an expression for x in terms of v . [5]
- 5 A particle P of mass $m\text{ kg}$ is attached to one end of a light elastic string of natural length 1.2 m and modulus of elasticity $0.75mg\text{ N}$. The other end of the string is attached to a fixed point O of a smooth plane inclined at 30° to the horizontal. P is released from rest at O and moves down the plane.
- (i) Show that the maximum speed of P is reached when the extension of the string is 0.8 m . [3]
- (ii) Find the maximum speed of P . [4]
- (iii) Find the maximum displacement of P from O . [4]

[Questions 6 and 7 are printed overleaf.]

6



A particle P of mass 0.4 kg is attached to one end of a light inextensible string of length 2 m . The other end of the string is attached to a fixed point O . With the string taut the particle is travelling in a circular path in a vertical plane. The angle between the string and the downward vertical is θ° (see diagram). When $\theta = 0$ the speed of P is 7 m s^{-1} .

(i) At the instant when the string is horizontal, find the speed of P and the tension in the string. [4]

(ii) At the instant when the string becomes slack, find the value of θ . [8]

7 A particle P , of mass $m\text{ kg}$, is attached to one end of a light elastic string of natural length 3.2 m and modulus of elasticity $4mg\text{ N}$. The other end of the string is attached to a fixed point A . The particle is released from rest at a point 4.8 m vertically below A . At time $t\text{ s}$ after P 's release P is $(4 + x)\text{ m}$ below A .

(i) Show that $4\frac{\text{d}^2x}{\text{d}t^2} = -49x$. [3]

P 's motion is simple harmonic.

(ii) Write down the amplitude of P 's motion and show that the string becomes slack instantaneously at intervals of approximately 1.8 s . [4]

A particle Q is attached to one end of a light **inextensible** string of length $L\text{ m}$. The other end of the string is attached to a fixed point B . The particle is released from rest with the string taut and inclined at a small angle with the downward vertical. At time $t\text{ s}$ after Q 's release BQ makes an angle of θ radians with the downward vertical.

(iii) Show that $\frac{\text{d}^2\theta}{\text{d}t^2} \approx -\frac{g}{L}\theta$. [3]

The period of the simple harmonic motion to which Q 's motion approximates is the same as the period of P 's motion.

(iv) Given that $\theta = 0.08$ when $t = 0$, find the speed of Q when $t = 0.25$. [5]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.