

**ADVANCED GCE
MATHEMATICS**

Further Pure Mathematics 3

4727

Candidates answer on the answer booklet.

OCR supplied materials:

- 8 page answer booklet (sent with general stationery)
- List of Formulae (MF1)

Other materials required:

- Scientific or graphical calculator

**Friday 28 January 2011
Morning**

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet. Please write clearly and in capital letters.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a scientific or graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- **You are reminded of the need for clear presentation in your answers.**
- The total number of marks for this paper is **72**.
- This document consists of **4** pages. Any blank pages are indicated.

- 1 (i) Find the general solution of the differential equation

$$\frac{dy}{dx} + xy = xe^{\frac{1}{2}x^2},$$

giving your answer in the form $y = f(x)$.

[4]

- (ii) Find the particular solution for which $y = 1$ when $x = 0$.

[2]

- 2 Two intersecting lines, lying in a plane p , have equations

$$\frac{x-1}{2} = \frac{y-3}{1} = \frac{z-4}{-3} \quad \text{and} \quad \frac{x-1}{-1} = \frac{y-3}{2} = \frac{z-4}{4}.$$

- (i) Obtain the equation of p in the form $2x - y + z = 3$.

[3]

- (ii) Plane q has equation $2x - y + z = 21$. Find the distance between p and q .

[3]

- 3 (i) Express $\sin \theta$ in terms of $e^{i\theta}$ and $e^{-i\theta}$ and show that

$$\sin^4 \theta \equiv \frac{1}{8}(\cos 4\theta - 4 \cos 2\theta + 3).$$

[4]


- (ii) Hence find the exact value of $\int_0^{\frac{1}{6}\pi} \sin^4 \theta \, d\theta$.

[4]

- 4 The cube roots of 1 are denoted by 1, ω and ω^2 , where the imaginary part of ω is positive.

- (i) Show that $1 + \omega + \omega^2 = 0$.

[2]



In the diagram, ABC is an equilateral triangle, labelled anticlockwise. The points A , B and C represent the complex numbers z_1 , z_2 and z_3 respectively.

- (ii) State the geometrical effect of multiplication by ω and hence explain why $z_1 - z_3 = \omega(z_3 - z_2)$.

[4]

- (iii) Hence show that $z_1 + \omega z_2 + \omega^2 z_3 = 0$.

[2]

- 5 (i) Find the general solution of the differential equation

$$3\frac{d^2y}{dx^2} + 5\frac{dy}{dx} - 2y = -2x + 13. \quad [7]$$

- (ii) Find the particular solution for which $y = -\frac{7}{2}$ and $\frac{dy}{dx} = 0$ when $x = 0$. [5]

- (iii) Write down the function to which y approximates when x is large and positive. [1]

- 6 Q is a multiplicative group of order 12.

- (i) Two elements of Q are a and r . It is given that r has order 6 and that $a^2 = r^3$. Find the orders of the elements a , a^2 , a^3 and r^2 . [4]

The table below shows the number of elements of Q with each possible order.

Order of element	1	2	3	4	6
Number of elements	1	1	2	6	2

G and H are the non-cyclic groups of order 4 and 6 respectively.

- (ii) Construct two tables, similar to the one above, to show the number of elements with each possible order for the groups G and H . Hence explain why there are no non-cyclic proper subgroups of Q . [5]

- 7 Three planes Π_1 , Π_2 and Π_3 have equations

$$\mathbf{r} \cdot (\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = 5, \quad \mathbf{r} \cdot (\mathbf{i} - \mathbf{j} + 3\mathbf{k}) = 6, \quad \mathbf{r} \cdot (\mathbf{i} + 5\mathbf{j} - 12\mathbf{k}) = 12,$$

respectively. Planes Π_1 and Π_2 intersect in a line l ; planes Π_2 and Π_3 intersect in a line m .

- (i) Show that l and m are in the same direction. [5]
- (ii) Write down what you can deduce about the line of intersection of planes Π_1 and Π_3 . [1]
- (iii) By considering the cartesian equations of Π_1 , Π_2 and Π_3 , or otherwise, determine whether or not the three planes have a common line of intersection. [4]

[Question 8 is printed overleaf.]

8 The operation $*$ is defined on the elements (x, y) , where $x, y \in \mathbb{R}$, by

$$(a, b) * (c, d) = (ac, ad + b).$$

It is given that the identity element is $(1, 0)$.

(i) Prove that $*$ is associative. [3]

(ii) Find all the elements which commute with $(1, 1)$. [3]

(iii) It is given that the particular element (m, n) has an inverse denoted by (p, q) , where

$$(m, n) * (p, q) = (p, q) * (m, n) = (1, 0).$$

Find (p, q) in terms of m and n . [2]

(iv) Find all self-inverse elements. [3]

(v) Give a reason why the elements (x, y) , under the operation $*$, do not form a group. [1]

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.