

# A Level Biology B (Advancing Biology)

H422/03 Practical skills in biology

# Monday 26 June 2017 – Morning

Time allowed: 1 hour 30 minutes

#### You must have:

- the Insert (inserted)
- a ruler (cm/mm)

#### You may use:

· a scientific or graphical calculator



First name	
Last name	
Centre number	Candidate number

#### **INSTRUCTIONS**

- The Insert will be found inside this document.
- Use black ink. You may use an HB pencil for graphs and diagrams.
- Complete the boxes above with your name, centre number and candidate number.
- Answer all the questions.
- Write your answer to each question in the space provided. If additional space is required, use the lined page(s) at the end of this booklet. The question number(s) must be clearly shown.
- Do **not** write in the barcodes.


### **INFORMATION**

- The total mark for this paper is 60.
- The marks for each question are shown in brackets [ ].
- Quality of extended responses will be assessed in questions marked with an asterisk (\*).
- · This document consists of 16 pages.



## Answer all the questions.

1 Fig. 1 is a fetal growth chart showing how biparietal diameter (BPD) increases during gestation.



- o 95th percentile
- × 50th percentile
- 5th percentile

Fig. 1

(a) Calculate the rate of growth at 30 weeks' gestation for the data shown in the uppermost curve (95th percentile). Show your working.

Answer =	 [2]	١

(b)	Three sets of data are shown in Fig. 1.
	How can the <b>three sets of data</b> be used together to monitor the health of a growing fetus?
	[1]
(c)	The measurements for BPD in Fig. 1 were taken using ultrasound.
	Evaluate the usefulness of ultrasound in measuring fetal growth.
	[2]

2	(a)*	Lettuce seeds r	equire a	minimum	day	length	to st	timulate	germina	ion.	Studies	have	shown
		that treatment v	with gibbe	erellins ca	an sti	mulate	gerr	mination	without	the r	need for	expos	sure to
		light.											

Gibberellic acid is a type of gibberellin.

A gardener wishes to extend the growing season for lettuces and wants to find out the minimum concentration of gibberellic acid to use to achieve germination.

Outline a practical method to determine the minimum concentration of gibberellic acid required to stimulate germination of lettuce seeds.

Your method should be based on the assumption that you are provided with the following:

•	school or college laboratory resources a solution of 100 mg dm <sup>-3</sup> gibberellic acid.
	ro.

- **(b)** A student is carrying out an experiment to determine the effect of a gibberellin, GA<sub>1</sub>, on the length of radicles (embryonic roots) of germinating pea seeds.
  - Fig. 2, on the insert, is a photograph of the germinating pea seeds.

All the seeds were soaked on the same day and have been growing for 5 days. The seeds have been soaked in different concentrations of  $GA_4$  as labelled in Fig. 2.

Using a ruler, measure the radicles in Fig. 2.

In the space below construct an appropriate table and record:

- the raw data to the most appropriate level of precision for this apparatus
- the mean values.

© OCR 2017 Turn over

[4]

3 A researcher investigated the effect of water temperature on the metabolic rate of a fish species.

For this investigation, the researcher used a type of respirometer attached to data-logging equipment containing an electronic sensor.

The electronic sensor measured any changes in the oxygen concentration of the water, which were used to calculate the metabolic rate of the fish.

Each of the following was included in the procedure for this investigation:

- Ten individual fish of the same species were used.
- The rate of oxygen consumption of each fish was measured.
- The temperature of the water was changed and the measurements repeated.
- After each set of measurements, the respirometers were flushed (cleaned and rinsed with hot water).

The rate of oxygen consumption was also measured in clean, empty respirometers. These measurements of oxygen consumption were too small to affect the outcome of the experiment.

(a) (i) The oxygen concentration of water can be measured using a number of different

	methods.	
	Give <b>one</b> advantage of using an electronic sensor with a data-logger.	
		. [1]
(ii)	State <b>two</b> of the control variables necessary in this experiment.	
		. [2]
(iii)	Suggest why the respirometers were flushed after each experiment.	
		-41

**(b)** Table 3.1 shows some of the results from this investigation.

Temperature of water in respirometer (°C)	Mean metabolic rate (mg O <sub>2</sub> kg <sup>-1</sup> hour <sup>-1</sup> )
10	86.0
16	131.6

Table 3.1

The standard deviation, *s*, for both means is 8.3.

A Student *t*-test can be performed to compare the mean metabolic rate at each temperature and determine if the two are significantly different from each other.

(i)	State the null hypothesis for this test.
	[1]
(ii)	Calculate the variance $(s^2)$ for these means.
	Answer =[1]

© OCR 2017 Turn over

(iii)	Using the formula	a below.	, calculate th	ne t value	for the	data in	Table 3.1.
-------	-------------------	----------	----------------	------------	---------	---------	------------

$$t = \frac{\overline{x}_A - \overline{x}_B}{\sqrt{\frac{s_A^2}{n_A} + \frac{s_B^2}{n_B}}}$$

Show your working. Give your answer to three decimal places.

<i>t</i> =	3.
ι –	 J

(iv) Using Table 3.2 and your answer to (b)(iii), comment on the mean metabolic rates for 10 °C and 16 °C.

Degrees of	Level of probability			
freedom	0.05	0.01	0.001	
8	2.306	3.355	5.041	
10	2.228	3.169	4.587	
18	2.101	2.878	3.922	
20	2.086	2.845	3.850	

Table 3.2

	 •	• • • • • • • • • • • • • • • • • • • •	 	
[3]				

(c)	The background research for this investigation into the effect of temperature on the metabolic rates of fish included information about increasing surface sea temperatures.
	It has been predicted that surface sea temperatures will continue to increase throughout the 21st century.
	Suggest <b>two</b> limitations of trying to link the results of this laboratory investigation with the likely consequences on marine fish as surface sea temperatures increase.
	[2]

© OCR 2017 Turn over

- An investigation into the effects of ageing on reaction times was carried out using data from a longitudinal study of the same people over a period of time.
  - A group of 25 people volunteered for a study in 1962.
  - As part of the study, the mean reaction time for each person was measured and recorded.
  - In 2012, ten of the same individuals were traced and volunteered to take part in a second study.
  - The mean reaction time using the same method was, again, measured and recorded.

Table 4 shows the results of the investigation.

Volunteer	Reaction time (10 <sup>-3</sup> s)		
	aged 18-25	aged 68–75	
1	262	310	
2	261	305	
3	262	298	
4	263	298	
5	262	297	
6	262	303	
7	258	311	
8	288	298	
9	290	300	
10	295	298	
Mean	270	302	
Median			
Range			

Table 4

(a) (i	i)	Complete Table 4 by calculating the median <b>and</b> the range for both data sets.	[2]
(ii	•	Evaluate the usefulness of the mean <b>and</b> the median when carrying out statistical analyses.	
			[2]
(iii	i)	Suggest <b>one</b> limitation of using the <b>range</b> in a statistical analysis.	
			[1]

(b)	The investigators decided to perform a paired <i>t</i> -test on the data in Table 4.
	Explain why a paired $t$ -test is more appropriate than an unpaired (Student's) $t$ -test for these data.
	[2]
(c)*	Laboratory experiments are often carried out to determine the effect of drugs on reaction times.
	These experiments may be:
	longitudinal studies of the same animal species over time
	carried out using human subjects
	carried out using different animal species.
	Discuss the ethical considerations of using humans <b>and</b> other animal species in laboratory experiments to determine the effect of drugs on reaction times.

**5** Researchers were studying the possible association between smoking and the respiratory tract infections, pneumonia and bronchitis.

The study included an analysis of the infection figures for **all** admissions to a hospital.

- Data from 5018 patients were collected.
- 4022 of these patients were **non-smokers**.
- Each patient had only one infection.

Some of the data from this study are shown in Table 5.1.

	Number of patients admitted with a respiratory tract infection		Number of patients	
	Pneumonia	Bronchitis	admitted with other type of infection	
Smokers	335	230	431	
Non-smokers	1166	526	2330	

Table 5.1

(a) Calculate the percentage of patients who were  ${\bf smokers}$ .

Give your answer to **two** decimal places.

		% [2]
(b)	The	researchers stated the following null hypothesis:
		Smoking does not affect the incidence of respiratory tract infections.
	(i)	Analyse the data in this study <b>and</b> comment on the researchers' hypothesis.
		In your answer you should use data to support your analysis.
		[3]
	(ii)	Suggest one statistical test that could be used to analyse the data further.

© OCR 2017

	(iii) Why would this statistical test be the most useful?
	[1]
(c)	Fig. 5.1, <b>on the insert</b> , is a photomicrograph showing tissues in the respiratory tract from one of the patients in the study.
	Name one structure, visible in Fig. 5.1, involved in the primary defence system.
	[1]
(d)	A sample of bacteria from a patient with pneumonia was collected and a pure culture was prepared.
	A Kirby-Bauer test was carried out to allow the doctor treating the patient to decide on the correct concentration of antibiotic to use.
	Six different concentrations of the antibiotic were used, as shown in Table 5.2.

Kirby-Bauer disc	Concentration of antibiotic (a.u.)
Α	0.0001
В	0.0010
С	0.0100
D	0.1000
E	1.0000
F	10.0000

Table 5.2

Describe the apparatus and method necessary to produce the six different concentrations.
You are supplied with an antibiotic solution of 10.0000 a.u. concentration and distilled water.
[4]

© OCR 2017 Turn over

(e) Fig. 5.2 shows the result of the Kirby-Bauer test.




Fig. 5.2

Use the information in Fig. 5.2 to advise the doctor on the most appropriate concentration of antibiotic to use. Explain your choice.
[2]

## **END OF QUESTION PAPER**

## 15

## **ADDITIONAL ANSWER SPACE**

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).			




#### Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

 $For queries \ or \ further \ information \ please \ contact \ the \ Copyright \ Team, \ First \ Floor, 9 \ Hills \ Road, \ Cambridge \ CB2 \ 1GE.$ 

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.