

Physics A

Advanced GCE **G484**

The Newtonian World

Mark Scheme for June 2010

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2010

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

Question	Expected Answers	Marks	Additional guidance
1(a)	The magnitude of the impulse on each object is the same Total energy is conserved	B1 B1	For 3 or 4 ticks mark and deduct 1 mark for each error.
(b) (i)	Correct use of $\frac{1}{2}mv^2$ Loss of KE = $0.03(144-81) = \mathbf{1.9}$ (or 1.89) J	C1 A1	0.27 J scores 1 st mark Do not allow 1.8
(b) (ii)	Change in momentum = $(0.06 \times 12) + (0.06 \times 9) = 1.26$ (Ns) Average force = rate of change of momentum = $1.26/0.15 = \mathbf{8.4}$ (or 8) N	C1 A1	Award 1 mark for 1.2 N ignore minus signs
(b) (iii)	8.4 N (or - 8.4)	B1	Allow ecf from (ii)
(c) (i)	ANY 3 of the following particles move with <u>rapid, random</u> motion (WTTE) elastic collisions negligible (or zero) volume of atoms (compared with volume of container) no intermolecular forces (except during collisions)/all internal energy is KE collision time negligible (compared to time between collision).	B1 B1 B1	Allow “gravitational force on molecules is negligible” Do not allow a bare “large number of particles”.
(c) (ii)	molecules make <u>collisions with walls/surface</u> (WTTE) (hence) exerts a force on the wall (or each collision has a change of momentum) Pressure = force/area	B1 B1 B1	Do not allow a bare “molecules collide with each other”
	Total	13	

Question	Expected Answers	Marks	Additional guidance
2 (a) (i)	Horizontal <u>component</u> of L provides the centripetal force (WTTE) Vertical <u>component</u> of L balances the weight (WTTE)	B1 B1	
(a) (ii)	$F = mv^2/r$ correct rearranged into $v = \sqrt{(Fr/m)}$ $v = \sqrt{(1.8 \times 10^6 \times 2000 / 1.2 \times 10^5)} = \mathbf{173 \text{ m s}^{-1}}$ (or 170)	C1 A1	Allow correct substitution of values into $F = mv^2/r$ for C1 mark
(b)	$mv^2/r = GMm/r^2$ $T = 2\pi r/v$ Correct manipulation of equations to give $T^2 = \frac{4\pi^2 r^3}{GM}$	B1 M1 A1	Do not allow a bare $v^2 = GM/r$ for the first mark – we need to see where this has come from.
(c) (i)	Equatorial orbit (WTTE) (<u>QWC</u> mark) Period is 24h/1day/same as Earth OR moves from West to East (WTTE)	B1 B1	QWC <u>equatorial</u> or <u>equator</u> must be spelled correctly
(c) (ii)	Correct rearrangement of $T^2 = (4\pi^2 r^3 / GM)$ to give $r^3 = T^2 GM / 4\pi^2$ correct sub. $r^3 = \{6.67 \times 10^{-11} \times 6.0 \times 10^{24} \times (8.64 \times 10^4)^2\} / 4\pi^2 = 7.57 \times 10^{22}$ $r = \mathbf{4.23 \times 10^7 \text{ m}}$ (or 4.2 or 4.3×10^7)	C1 C1 A1	(1 day = 8.64×10^4 s is given on the data sheet). For those who use $g = GM/r^2$ with $g = 9.81$ award 1 mark for $r = 6.4 \times 10^6$ m.
	Total	12	

Question	Expected Answers	Marks	Additional guidance
3(a)	Acceleration is (directly) proportional to the displacement/distance (from the equilibrium position/central pt) Acceleration is always directed towards the equilibrium position/central point.	B1 B1	Allow “fixed point” or “point” Allow acc. is in opposite direction to displacement (WTTE) If formula is used: allow $a \propto -x$ for 1 st mark and 2 nd mark if x is stated as displacement.
(b) (i)	Curve symmetrical about energy axis with maximum at 18 zero at +0.04 and – 0.04	B1 B1	Ignore points where graphs cross Give bod if not labelled K but correct
(b) (ii)	Horizontal straight line passing 18	B1	Give bod if not labelled T but correct
(c) (i)	0.04 m	B1	
(c) (ii)	$\frac{1}{2}m(v_{\max})^2 = 0.018$ $v_{\max} = \sqrt{(2 \times 0.018 / 0.12)} = \mathbf{0.55} \text{ ms}^{-1}$ (0.548)	C1 A1	Many will use 18 instead of 0.018. This results in 17.3 and scores 1 mark. Allow ecf for cand's value of max KE. Do not allow 0.54 for second mark.
(c) (iii)	correct use of $v_{\max} = 2\pi fA$ $f = (0.55 / 0.04 \times 2\pi) = \mathbf{2.2}$ (or 2.19 or 2.18)Hz	C1 A1	Allow ecf for cand's values from (c)(i) and/or (c) (ii). E.g for 17.3 $f = 68.8$ Hz. This scores 2 marks e.c.f. Do not allow 2.1
(d)	Award first mark for stating the ‘ driver ’ of the oscillations and the second mark for stating what is ‘ driven ’ i.e. oscillating useful applications: e.g. Cooking: micro waves cause water molecules to resonate Woodwind: reed causes air column to resonate Brass: lips cause air column to resonate MRI: radio waves (in a magnetic field) cause nuclei/proton to resonate Radios: radio waves cause electrons/current to resonate Person on swing: intermittent pushes cause swing to resonate problem: Bridges: wind/walkers causes bridge to resonate Vehicles: engine vibrations cause panels/mirrors to resonate Earthquakes: ground vibrating causes buildings to resonate	B1 B1 B1 B1	No marks to be awarded for a bare statement of the example e.g MRI. Please allow any other valid examples.
		Total	14

Question	Expected Answers	Marks	Additional guidance
4 (a) (i)	Brownian (motion) (QWC mark)	B1	QWC <u>Brownian</u> spelled correctly
(a) (ii)	ANY two from the following three: air molecules are moving in different directions/randomly with different speeds mass/size of air molecules is smaller than smoke particles	B1 B1	Answers that refer to smoke particles only cannot score the marks.
(b) (i)	vol = $(4/3) \pi r^3 = 5.58 \times 10^{-3}$ correct sub into $pV = nRT$ i.e. with T as 290K $n = (2.6 \times 10^5 \times 5.58 \times 10^{-3}) / 8.31 \times 290 = 0.602$ moles mass = $n \times 0.028 = \mathbf{0.0169}$ kg (0.016856)	C1 C1 A1 A1	Allow ecf for wrong volume Allow use of $pV = NkT$ and $n = N/N_A$ Allow ecf for cand's value for n If 17° C used allow maximum of 2 marks for $n = 10.3$ moles and $m = 0.29$ kg
(b) (ii) 1	no net heat flow between objects (WTTE)	B1	Allow "they are at the same temp."
(b) (ii) 2	correct use of $P/T = \text{constant}$: e.g. $P = (273/290) \times 2.6 \times 10^5$ $P = \mathbf{2.45 \times 10^5}$ (or 2.4×10^5 or 2.5×10^5) Pa	C1 A1	Allow correct use of $pV = nRT$
Total		10	

Question	Expected Answers	Marks	Additional guidance
5(a) (i)	Initial KE of car = $0.5 \times 970 \times 27^2 = \mathbf{3.5 \times 10^5 \text{ J}}$ (353565J)	B1	
(a) (ii)	Work done = Av Force x distance moved Av Force = $3.5 \times 10^5 \text{ J} / 40 = \mathbf{8.8 \times 10^3 \text{ N}}$ (or 8750 N) (or $353565 / 40 = 8836.7 \text{ N}$) Assumption: no air resistance	C1 A1 B1	If $v^2 = u^2 + 2as$ is used. accept $a = 0 - 27^2 / (2 \times 40) = 9.113 \text{ ms}^{-2}$ C1 $F = ma = 970 \times 9.11 = 8.84 \times 10^3 \text{ N}$ A1 Allow air friction or drag
(b) (i)	correct use of $E = mc\Delta\theta$: $3.5 \times 10^5 / 4 = 1.2 \times 520 \times \Delta\theta$ $\Delta\theta = \mathbf{140^\circ\text{C}}$ (if 353565 is used $\Delta\theta = 142^\circ\text{C}$)	C1 A1	If cand. forgets to divide by 4 allow any value between 560 and 570 for 1 mark.
(b) (ii)	<u>Air resistance</u> will be acting (slowing down the car) (hence) <u>reducing the KE of the car</u> (WTTE) The <u>discs are hotter</u> than the surroundings (hence) <u>energy/heat</u> will be lost from <u>discs/brakes</u> (WTTE)	M1 A1 B1 B1	Do not allow sound since only a tiny proportion of energy is lost in this way. Allow other valid comments as alternative ways of scoring one or both of the 'B' marks: e.g. 'hot spots' on discs; discs are different. Try to credit a well argued case based upon correct physics- e.g. wheels locking.
(b) (iii)	Any valid suggestion: e.g. use a material with a higher s.h.c use a disc with a higher heat capacity Use discs of greater mass put holes in the discs (to increase air flow)	B1	Confusion between shc and heat capacity should not be penalised.
Total		11	

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2010

