

Monday 13 May 2019 – Afternoon

AS Level Further Mathematics A

Y531/01 Pure Core

Time allowed: 1 hour 15 minutes



You must have:

- Printed Answer Booklet
- Formulae AS Level Further Mathematics A

You may use:

• a scientific or graphical calculator

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- Write your answer to each question in the space provided in the Printed Answer Booklet. If additional space is required, you should use the lined page(s) at the end of the Printed Answer Booklet. The question number(s) must be clearly shown.
- You are permitted to use a scientific or graphical calculator in this paper.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question.
- The acceleration due to gravity is denoted by $gm s^{-2}$. Unless otherwise instructed, when a numerical value is needed, use g = 9.8.

INFORMATION

- The total mark for this paper is **60**.
- The marks for each question are shown in brackets [].
- You are reminded of the need for clear presentation in your answers.
- The Printed Answer Booklet consists of 12 pages. The Question Paper consists of 4 pages.

Answer all the questions.

- 1 You are given that z = 3 4i.
 - (a) Find
 - z,
 - $\arg(z)$,

On an Argand diagram the complex number w is represented by the point A and w^* is represented by the point B.

- (b) Describe the geometrical relationship between the points *A* and *B*. [2]
- 2 Matrices **P** and **Q** are given by $\mathbf{P} = \begin{pmatrix} 1 & k & 0 \\ -2 & 1 & 3 \end{pmatrix}$ and $\mathbf{Q} = ((1+k) 1)$ where k is a constant.

Exactly one of statements A and B is true.

Statement A:	P and Q (in that order) are conformable for multiplication.
Statement B:	Q and P (in that order) are conformable for multiplication.

- (a) State, with a reason, which one of A and B is true. [2]
- (b) Find either PQ or QP in terms of k. [2]
- 3 The position vector of point A is $\mathbf{a} = -9\mathbf{i} + 2\mathbf{j} + 6\mathbf{k}$. The line l passes through A and is perpendicular to \mathbf{a} .

(a) Determine the shortest distance between the origin, <i>O</i> , and <i>l</i> .	[2]
<i>l</i> is also perpendicular to the vector b where $\mathbf{b} = -2\mathbf{i} + \mathbf{j} + \mathbf{k}$.	
(b) Find a vector which is perpendicular to both a and b .	[1]
(c) Write down an equation of <i>l</i> in vector form.	[1]
<i>P</i> is a point on <i>l</i> such that $PA = 2OA$.	

(d) Find angle *POA* giving your answer to 3 significant figures. [3]

C is a point whose position vector, **c**, is given by $\mathbf{c} = p\mathbf{a}$ for some constant *p*. The line *m* passes through *C* and has equation $\mathbf{r} = \mathbf{c} + \mu \mathbf{b}$. The point with position vector $9\mathbf{i} + 8\mathbf{j} - 12\mathbf{k}$ lies on *m*.

(e) Find the value of p. [3]

4 In this question you must show detailed reasoning.

You are given that $f(z) = 4z^4 - 12z^3 + 41z^2 - 128z + 185$ and that 2+i is a root of the equation f(z) = 0.

(a) Express f(z) as the product of two quadratic factors with integer coefficients. [5]

(b) Solve
$$f(z) = 0$$
. [3]

Two loci on an Argand diagram are defined by $C_1 = \{z : |z| = r_1\}$ and $C_2 = \{z : |z| = r_2\}$ where $r_1 > r_2$. You are given that two of the points representing the roots of f(z) = 0 are on C_1 and two are on C_2 . *R* is the region on the Argand diagram between C_1 and C_2 .

- (c) Find the exact area of *R*. [4]
- (d) ω is the sum of all the roots of f(z) = 0.

Determine whether or not the point on the Argand diagram which represents ω lies in R. [2]

5 In this question you must show detailed reasoning.

You are given that α , β and γ are the roots of the equation $5x^3 - 2x^2 + 3x + 1 = 0$.

- (a) Find the value of $\alpha^2 \beta^2 + \beta^2 \gamma^2 + \gamma^2 \alpha^2$. [5]
- (b) Find a cubic equation whose roots are α^2 , β^2 and γ^2 giving your answer in the form $ax^3 + bx^2 + cx + d = 0$ where a, b, c and d are integers. [4]
- 6 A transformation T is represented by the matrix T where $T = \begin{pmatrix} x^2 + 1 & -4 \\ 3 2x^2 & x^2 + 5 \end{pmatrix}$. A quadrilateral Q, whose area is 12 units, is transformed by T to Q'.

Find the smallest possible value of the area of Q'.

Turn over for questions 7 and 8

[5]

7 A transformation A is represented by the matrix A where $\mathbf{A} = \begin{pmatrix} -1 & x & 2 \\ 7-x & -6 & 1 \\ 5 & -5x & 2x \end{pmatrix}$.

The tetrahedron H has vertices at O, P, Q and R. The volume of H is 6 units.

- P', Q', R' and H' are the images of P, Q, R and H under A.
- (a) In the case where x = 5
 - find the volume of H',
 - determine whether A preserves the orientation of *H*. [3]
- (b) Find the values of x for which O, P', Q' and R' are coplanar (i.e. the four points lie in the same plane).[4]

8 In this question you must show detailed reasoning.

M is the matrix $\begin{pmatrix} 1 & 6 \\ 0 & 2 \end{pmatrix}$. Prove that $\mathbf{M}^n = \begin{pmatrix} 1 & 3(2^{n+1}-2) \\ 0 & 2^n \end{pmatrix}$, for any positive integer *n*. [6]

END OF QUESTION PAPER



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.