

# F

## Friday 27 May 2022 – Morning

## GCSE (9–1) Combined Science B (Twenty First Century Science)

J260/02 Chemistry (Foundation Tier)

Time allowed: 1 hour 45 minutes

#### You must have:

- a ruler (cm/mm)
- the Data Sheet for GCSE (9–1) Combined Science (Chemistry) B (inside this document)

#### You can use:

- an HB pencil
- · a scientific or graphical calculator



Please write clear	ly in black i	nk. <b>Do not v</b>	vrite in the barcodes.		
Centre number			Candidate number		
First name(s)					
Last name _					

#### **INSTRUCTIONS**

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.
- Answer all the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

#### **INFORMATION**

- The total mark for this paper is 95.
- The marks for each question are shown in brackets [].
- Quality of extended response will be assessed in questions marked with an asterisk (\*).
- This document has 28 pages.

#### **ADVICE**

Read each question carefully before you start your answer.



#### Answer **all** the questions.

1 Cold packs are used to treat some sports injuries.

They contain ammonium nitrate and water.

- (a) The formula of ammonium nitrate is NH<sub>4</sub>NO<sub>3</sub>.
  - (i) Name the **three** elements present in ammonium nitrate.

1																								
ı																							 	


[3]

(ii) What is the total number of atoms shown in the formula?


[1]

**(b)** When the ammonium nitrate and water mix in the cold pack, the temperature of the mix decreases.

The diagram shows the reaction profile for this reaction.



Progress of reaction

(i) What is the name given to a change which makes the temperature of the mix decrease?

Put a (ring) around the correct option.

Combustion Endothermic Exothermic Freezing

[1]

(ii) Draw lines to connect each label with its correct position on the diagram.

Label	Position	
Activation energy	A	
Energy of reaction	В	
Products	С	
Reactants	D	
		[3]

**(c)** Kofi plans an experiment to find the energy change when the ammonium nitrate from a cold pack dissolves in water.

The energy change depends on the mass of water and the temperature change.

What apparatus will he need to take the measurements?

Balance
Burette
Gas syringe
pH meter
Pipette

Thermometer

Tick (✓) **two** boxes.

[2]

(d) Here are Kofi's res
-------------------------

Mass of water used = 250 g

Temperature of water before adding ammonium nitrate = 20.2 °C

Temperature of water after adding ammonium nitrate = 4.8 °C

Calculate the energy change when the ammonium nitrate from the cold pack dissolves in water.

Use the formula: Energy change (J) =  $4.2 \times \text{mass}$  of water (g)  $\times \text{temperature change}$  (°C)

Give your answer to 3 significant figures.

Energy change = ...... J [3]

## 5 BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE


- ${\bf 2} \qquad {\rm Harmful\ gases\ such\ as\ NO,\ CO\ and\ SO}_2\ are\ emitted\ from\ car\ engines\ running\ on\ fossil\ fuels.}$ 
  - (a) These gases are produced in the car engine.

Draw lines to connect each gas with its formula and how it is formed.

Gas	Formula	How it is formed
Sulfur dioxide	СО	Combustion of sulfur impurities in the fuel
Carbon monoxide	NO	From air at high temperature
Nitrogen monoxide	SO <sub>2</sub>	Incomplete combustion of the fuel

[4]

**(b)** Some harmful gases are removed by a catalytic converter before leaving the exhaust pipe of the car. The diagram shows gases passing through a catalytic converter coated with a catalyst.



The harmful gases from the car engine do not react with each other until they are in contact with the hot catalyst.

(i)	Why do the gases not react with each other until they are in contact with the hot
	catalyst?

Tick (✓) one box.

The reaction has a high activation energy.

The reaction is exothermic.

The reaction is too fast.

The catalyst is used up in the reaction.

(c) The equation shows one reaction of the gases in the catalytic converter.

2CO + 2NO 
$$\rightarrow$$
 2CO<sub>2</sub> + N<sub>2</sub>

Identify which gas is oxidised.

Explain your answer.

Gas which is oxidised.....

Explanation .....


[1]

2 1.	10anonium	budrovida	rooto wi	h h	ماحم مامام حنام	aaid in	a nautraliaatian	roostion
<b>ວ</b> Ⅳ	naunesium	nvaroxiae	reacts wi	11 1170	arocnionc	acid in	a neutralisation	reaction.

(a)	(i)	Complete the balanced symbol equation for the reaction between magnesium hydroxide
		and hydrochloric acid.

$$\label{eq:mgOH} \mathrm{Mg(OH)}_2 \ + \ .....\mathrm{HC} l \ \rightarrow \ \mathrm{MgC} l_2 \ + \ .....\mathrm{H}_2\mathrm{O} \hspace{1cm} \text{[1]}$$

(ii) Draw lines to connect each compound from the reaction with its correct description.



(b) Complete the sentence about neutralisation.

Use the symbols and words from the list.

acid	alkali	water	C1 <sup>-</sup>	H⁺	Mg <sup>2+</sup>	OH-
Neutralisat	tion occurs whe	en		ions fr	om an	
		react with	l		ions fro	m an
		to form				[3

(c) Milk of magnesia is a medicine used to neutralise ex	excess acid in the stomach	
--	----------------------------	--

Milk of Magnesia	
One 15 cm <sup>3</sup> spoonful contains 1200 mg of magnesium hydroxide.	

Leo does a titration with hydrochloric acid to check the amount of magnesium hydroxide in one spoonful of Milk of Magnesia.

This is the method:

- Add 15 cm<sup>3</sup> of Milk of Magnesia into a conical flask
- Add a few drops of indicator
- Add hydrochloric acid from a measuring cylinder until all of the magnesium hydroxide is neutralised.

When should Leo stop adding the acid?

lick (✓) one box.	
When all the acid has been added.	
When he has added 15 cm <sup>3</sup> of acid.	
When the conical flask is full.	
When the indicator changes colour.	

© OCR 2022 Turn over

[1]

(d)	) Leo decides to measure the acid with a more accurate piece of apparatus instead of the measuring cylinder.					Э			
	Which piece of apparatus should he use?								
	Put	a (ring) aro	und the correc	t answer.					
	Bea	ıker	Burette	Flask	The	rmometer			[1]
(e)	Leo	repeats the	e titration with t	the more ac	curate piec	e of appara	tus.		
	Her	e are his re	sults.						
				1	2	3	4	5	
	Vo	lume of aci	d used (cm <sup>3</sup> )	29.30	27.20	27.30	27.20	27.30	
	Mean volume of acid used =					ōcm³ spoon			
		Use the fo	rmula:						
		Mass of m	agnesium hyd	roxide(g) =	volume of h	ydrochloric	acid(cm³) ×	0.044	
			Volume	of hydroch	loric acid =			CI	m³ <b>[3]</b>
			Volatile	. 51 11, 410011	iono dola				[∿]

## 11 BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

- 4 Sodium is a metal and oxygen is a non-metal.
  - (a) Complete the table to describe the difference in the properties of metals and non-metals.Use words from the list.

dull high low s	iny
-----------------	-----

Property	Metals	Non-metals
Melting point	high	low
Boiling point		
Density		
Appearance		
Electrical conductivity	good	none

[3]

(b) Complete the sentence to explain why sodium is a good conductor of electricity.

Put a ring around the **two** correct answers.

Sodium is a good conductor of electricity because it has **electrons / ions / molecules** which are **fixed / mobile / static**. [2]

- (c) Sodium has an electronic structure 2.8.1 and oxygen has an electronic structure 2.6.
  - (i) Complete the table to show the position of these two elements in the Periodic Table.Use the Data Sheet.

	Period	Group
Sodium		
Oxygen		

[2]

(ii) Sodium and oxygen react together to form sodium oxide.

Complete the table to show the electronic structures and charges of the ions formed.

	Electronic structure	Charge
Sodium ion		
Oxide ion		

[3]

- 5 Paper chromatography can be used to identify dyes in a mixture.
  - (a) Complete the sentence to describe how chromatography works.

Use words from the list.

condensed	distributed	dyes	mobile	paper	solvent	stationary	y
The		in the	mixture are	separated v	when they ar	re	
	b	etween th	ıe		wh	nich is the	
	р	hase and	the			which is the	
	p	hase.					[4]

(b) Rf values are used to identify dyes. They are calculated using this equation:

 $Rf = \frac{\text{distance moved by the substance}}{\text{distance moved by the solvent}}$ 

The Rf value of a dye can be calculated from the result of an experiment and compared with Rf values in a data book.

- **Table 5.1** shows the results of an experiment to identify an unknown dye.
- **Table 5.2** shows the Rf values of some dyes in a data book.

Distance moved by the dye (cm)	3.5
Distance moved by the solvent (cm)	8.3

	Rf value
Marigold Orange	0.36
Mustard Yellow	0.89
Sunshine Yellow	0.42

Table 5.1 Table 5.2

Which dye has been identified from the experimental results in <b>Table 5.1</b> ?
Use a calculation to explain your answer.
rs

(c)\* A student checks the Rf value of sunshine yellow using chromatography.

The chromatogram produced by the student is shown for the sunshine yellow dye.





Fig. 5.1

Describe how the student did the experiment to find the Rf value of the dye.
Include the apparatus used and any measurements taken.
[6]

6 The diagrams show the structures of some forms of carbon.



(	a)	(i)	How do	the diagrams	show that al	I the structures	are the same	e element?
١	ч,	1''	11000 00	uio alagiailio	oriow triat ar	i ti io oti aotai oo	are the earne	ololliolit.

Tick (✓)	one	box.
----------	-----	------

There is only one type of atom in all of the structure	es.
--	-----

They are all the same shape.

They are all 3 dimensional.

They all have the same number of atoms.

[1]

(ii) Which statements about these structures are **true only for diamond**, which are **true only for fullerenes**, and which are **true for all the structures**?

Tick (✓) one box in each row.

	True only for diamond	True only for fullerenes	True for all the structures
Each carbon atom is bonded to 3 others.			
All atoms are held together by covalent bonds.			
Carbon atoms are arranged in a tetrahedral (pyramidal) shape.			

[3]

(b)	Full	erenes are nan	oparticles.			
	(i)	What is the approximate size of a nanoparticle?				
		Put a ring around the correct answer.				
		≤100 nm 1000 nm ≥10 000 nm				
	(ii)	Fullerenes can be used to carry drugs into the body.				[1]
		Which <b>two</b> sta	tements expla	in why?		
		Tick (✓) <b>two</b> b	oxes.			
		They are smal	l enough to pa	ss through body	tissues.	
	They are made of carbon atoms.					
	They can fit molecules inside them.  They are good for you.					
		They can act as catalysts.				
	(iii)			vith the use of na enefits outweigh	noparticles to carry drug the risks.	[2] s into the human
		Suggest <b>one</b> rinto the body.	isk and <b>one</b> be	enefit to a persor	n's health of using fullere	enes to carry drugs
		Risk				
		Benefit				
						[21

7 Table 7.1 shows some information about the elements in Group 17(7) of the Periodic Table.

	State and colour at 20 °C	Reaction with iron wool	Product
Fluorine	pale yellow gas	Reacts instantly.	Iron fluoride
Chlorine	yellow-green gas	Reacts with heated iron wool very quickly, although not as quickly as fluorine does.	Iron chloride
Bromine	deep red liquid	Has to be warmed and the iron wool heated. The reaction is faster than that of iodine but slower than that of chlorine.	Iron bromide
lodine	grey solid	Has to be heated strongly and so does the iron wool. The reaction is slow.	Iron iodide

#### Table 7.1

(a)	(i)	Name the most reactive member of Group 17(7).
		[1]
	(ii)	Name <b>one</b> member of Group 17(7) that is not shown in <b>Table 7.1</b> .
		[1]
(	(iii)	Predict the state and colour at 20 °C, the reaction with iron wool and the product of the reaction for the element identified in (ii).
		State and colour at 20 °C
		Reaction with iron wool
		Product formed with iron wool
		[3]

**(b)** The elements in Group 17(7) form compounds with other metals.

**Table 7.2** shows information about some of these compounds.

lons	Formula	Relative formula mass
	KF	58.1
Ca <sup>2+</sup> and Br <sup>-</sup>		199.9
Fe <sup>3+</sup> and F <sup>-</sup>	FeF <sub>3</sub>	

Table 7.2

(i)	Complete the table to show the missing ions and formula.	[2]
(ii)	Calculate the relative formula mass for FeF <sub>3</sub> .	
	Use the Data Sheet.	
	Polativo formula mass –	[2]
	Relative formula mass =	. [2

8 The table gives information about some of the compounds present in crude oil.

Number of carbon atoms	Molecular formula	Empirical formula	Melting point (°C)	Boiling point (°C)	State at room temperature
4	C <sub>4</sub> H <sub>10</sub>	C <sub>2</sub> H <sub>5</sub>	-138	0	Gas
5		C <sub>5</sub> H <sub>12</sub>	-130	36	
6	C <sub>6</sub> H <sub>14</sub>	C <sub>3</sub> H <sub>7</sub>	<b>–</b> 95	69	Liquid
7	C <sub>7</sub> H <sub>16</sub>	C <sub>7</sub> H <sub>16</sub>	-90		Liquid
8	C <sub>8</sub> H <sub>18</sub>		<b>–</b> 57	126	Liquid

(a)	(i)	Complete the table to show the missing molecular formula and empirical formula.	[2]
	(ii)	Predict the boiling point for the compound with 7 carbon atoms.	
		Boiling point =°	C [1]
	(iii)	Predict the state of the 5 carbon compound at room temperature (20 °C).	
		Explain your answer.	
		State	
		Explanation	
			[2]
			141

(b) All the compounds in the table are in the same homologous series.

All	members of a homologous series have the same general formula.
(i)	Give <b>two</b> other characteristics of a homologous series that are shown in the table.
	1
	2
	[2
(ii)	Complete the sentences to describe the compounds present in crude oil that are shown in the table.
	Put a ring around each correct answer.
	Crude oil is a mixture of hydrocarbons / polymers / salts.
	The compounds are from the homologous series allotropes / alkanes / alkenes.
	They all have the general formula $C_nH_{2n}$ / $C_nH_{2n+1}$ / $C_nH_{2n+2}$ .

[3]


**9** Solid calcium carbonate reacts with dilute hydrochloric acid to form calcium chloride, carbon dioxide and water.

$${\rm CaCO_3(s)} \ + \ 2{\rm HC}\mathit{l}({\rm aq}) \ \longrightarrow \ {\rm CaC}\mathit{l}_2({\rm aq}) \ + \ {\rm CO}_2({\rm g}) \ + \ {\rm H}_2{\rm O(I)}$$

(a) Jane investigates the rate of this reaction. She measures the change in mass during the reaction over five minutes.

She uses 10 g of calcium carbonate lumps and 50 cm<sup>3</sup> of dilute hydrochloric acid.

The graph shows Jane's results.



	(i)	What was the time taken for the reaction to finish?	
	(ii)	What was the total mass lost during the reaction?	[1]
	(iii)	Total mass lost =	[2]
		Rate of reaction = g/s	[2]
(b)		e repeats the experiment with 10g of calcium carbonate <b>powder</b> instead of 10g of ps. She keeps everything else the same.	
	Ske	etch a line on the graph to show the results she should expect.	[2]
(c)		nplete the sentences to explain why the rate of reaction changes when powdered sium carbonate is used instead of lumps.	
	Put	a ring around each correct answer.	
		e surface area of 10g of powdered calcium carbonate is <b>larger than / smaller than / thee as</b> 10g of lumps.	ne
		total volume of 10 g of powdered calcium carbonate is larger than / smaller than / th	e
	รสก	ne as 10g of lumps.	[2]

## **END OF QUESTION PAPER**

### 24

## **ADDITIONAL ANSWER SPACE**

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).					



•••••	

 ,	 	
 <u> </u>	 	



#### **Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.