

# Tuesday 13 June 2023 – Morning GCSE (9–1) Chemistry A (Gateway Science)

J248/04 Paper 4 (Higher Tier)

Time allowed: 1 hour 45 minutes

### You must have:

- a ruler (cm/mm)
- the Data Sheet for GCSE (9–1) Chemistry A (inside this document)

#### You can use:

- · a scientific or graphical calculator
- an HB pencil





Please write clea	rly in blac	k ink. <b>Do</b>	not wri	te in the barcodes.		
Centre number				Candidate number		
First name(s)						
Last name						

### **INSTRUCTIONS**


- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.
- · Answer all the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for a correct method, even if your answer is wrong.

## **INFORMATION**

- The total mark for this paper is 90.
- The marks for each question are shown in brackets [ ].
- Quality of extended response will be assessed in questions marked with an asterisk (\*).
- This document has 28 pages.

#### **ADVICE**

· Read each question carefully before you start your answer.



## **Section A**

You should spend a maximum of 30 minutes on this section.

Write your answer to each question in the box provided.

1 Crude oil is a resource that is being made extremely slowly.

Which word describes a resource that is being made extremely slowly?

- **A** Finite
- **B** Hydrocarbon
- C Non-renewable
- **D** Petrochemical

Your answer		[1]
-------------	--	-----

2 The Group 7 element fluorine is a gas at room temperature and pressure.

The Group 7 elements show a trend in boiling points going down the group.

Which row shows the boiling points of the Group 7 elements?

	Boiling Point (°C)								
	Fluorine	Chlorine	Bromine	lodine					
Α	-188	59	184	-15					
В	-188	-34	59	184					
С	188	184	59	<b>–15</b>					
D	<b>–</b> 15	184	188	59					

Your answer	[1]

3	Cop	Copper can be extracted from copper ore by heating the copper ore with carbon.							
	Cop	Copper can also be extracted by bioleaching.							
	What is an advantage of bioleaching?								
	Α	It can produce sulfuric acid.							
	В	It extracts copper, which is then purified by electrolysis.							
	С	It is done at low temperatures.							
	<b>D</b> It is slow.								
	Your answer								
4	Wh	/hich gas was the most abundant in the Earth's early atmosphere?							
	Α	Argon							
	В	Carbon dioxide							
	С	Nitrogen							
	D	Oxygen							
	Υοι	ur answer	[1]						

5	This is the	halanced s	vmbol equat	tion for the	reaction of	f conner (	oxide with	carbon
J	11115 15 1116	Dalaliceu S	viiibui <del>e</del> qua	uon ioi ine	Teachon of	i coppei (	JAIUE WILLI	carbon.

$$2CuO + C \rightarrow 2Cu + CO_2$$

What happens to the copper oxide in this reaction?

- **A** It is neutralised.
- **B** It is oxidised.
- **C** It is reduced.
- **D** It is thermally decomposed.


Your answer		[1]
-------------	--	-----

# 6 Which row in the table shows three greenhouse gases?

Α	argon	carbon dioxide	nitrogen
В	carbon dioxide	methane	water vapour
С	hydrogen	methane	water vapour
D	carbon dioxide	nitrogen	water vapour

Your answer	[1]

7 The graph shows the volume of hydrogen gas made in an experiment.



What is the rate of reaction when the time is 20s?

- **A**  $0.44 \, \text{cm}^3/\text{s}$
- **B**  $2.25\,\text{cm}^3/\text{s}$
- **C**  $25 \, \text{cm}^3 / \text{s}$
- **D**  $900 \, \text{cm}^3/\text{s}$

Your answer		[1]
-------------	--	-----

- **8** Why are alloys stronger than the pure metals which they are made from?
  - A Alloys combine the properties of the metals they are made from.
  - **B** Alloys contain atoms of different sizes.
  - **C** Alloys contain different atoms bonded together.
  - **D** Alloys have strong bonds between their molecules.

Your answer
Your answer

9	Whi	ich statement about chemical cells is correct?	
	Α	They produce a voltage indefinitely.	
	В	They produce a voltage once all of the reactants are used up.	
	С	They produce a voltage until one of the reactants is completely used up.	
	D	They produce a voltage until the reactants are partly used up.	
	You	er answer	[1]
10	Whi	ich of these homologous series can form addition polymers?	
	Α	Alcohols and carboxylic acids	
	В	Alkenes and alkanes	
	С	Alkenes only	
	D	Carboxylic acids only	
	You	er answer	[1]
11	Whi	ich statement about the greenhouse effect is correct?	
	Α	Greenhouse gases absorb all the infrared radiation that is emitted by the Earth's surface.	
	В	Greenhouse gases make up a large percentage of the Earth's current atmosphere.	
	С	The greenhouse effect is caused by the absorption and reflection of infrared radiation by greenhouse gases.	
	D	The higher the concentration of greenhouse gases in the Earth's atmosphere, the colder Earth is likely to become.	the
	You	r answer	[1]


**12** Ethane is a very small hydrocarbon molecule.


Which row about ethane is correct?


	Easy to ignite?	Boiling point
Α	✓ high	
В	X	high
С	✓	low
D	Х	low


Your answer [1]

13 Which graph shows a dynamic equilibrium?









Your answer

© OCR 2023 Turn over

[1]

**14** Ethene,  $C_2H_4$ , reacts with bromine in an addition reaction.


What is the structural formula of the product?

Α


В



C



D



Your answer

[1]

15 What is the half equation for the reaction at the anode in a hydrogen/oxygen fuel cell?

$$\mathbf{A} \quad 2\mathrm{H_2} + \mathrm{O_2} \rightarrow 2\mathrm{H_2O}$$

$$\mathbf{B} \quad 2\mathrm{H}_2 \rightarrow 4\mathrm{H}^+ + 4\mathrm{e}^-$$

$$\mathbf{C} \quad 4\mathrm{H^+} + 4\mathrm{e^-} \rightarrow 2\mathrm{H_2}$$

$$\mathbf{D} \quad 4\mathrm{H^+} + \mathrm{O_2} + 4\mathrm{e^-} \rightarrow 2\mathrm{H_2O}$$

Your answer

[1]

9

# **BLANK PAGE**

PLEASE DO NOT WRITE ON THIS PAGE

## **Section B**

16 A student investigates two solutions, A and B.

They know that

- one solution contains a halide ion
- the other solution contains a different anion.

They test 2 cm<sup>3</sup> of each solution for the halide ion using this method:

- Add a few drops of dilute hydrochloric acid and shake.
- Add a few drops of silver nitrate solution.
- Record the colour of the precipitate.
- (a) The teacher says that the student should have used a different acid instead of dilute hydrochloric acid.

(i)	State the name of the acid the student should have used.	
		[1]
(ii)	Explain why using dilute hydrochloric acid would affect the results of this test.	
		[2

(b) The student repeats the test for halide ions using the correct acid and silver nitrate solution.

They also test each solution using a few drops of dilute hydrochloric acid followed by a few drops of barium chloride solution.

The table shows their results.

Solution	Observation with silver nitrate solution	Observation with barium chloride solution	
Α	cream precipitate	no change	
В	no change	white precipitate	

State the name of the anion in each solution.	
Solution A	
Solution <b>B</b>	[2]

	11
(c)	Solution <b>A</b> also contains copper ions, Cu <sup>2+</sup> .
	Copper ions react with hydroxide ions, OH <sup>-</sup> , to make a precipitate of copper(II) hydroxide.
	Write the <b>balanced ionic</b> equation for this reaction. Include the state symbols.
	[3]
(d)	Barium chloride solid is toxic if swallowed and harmful if inhaled.
	Barium chloride solutions with concentrations of $21\mathrm{g/dm^3}$ are suitable for experiments in school.
	Calculate the mass of barium chloride that should be dissolved in $25\mathrm{cm}^3$ of water to make a solution with a concentration of $21\mathrm{g}/\mathrm{dm}^3$ .
	Use the equation:
	$concentration = \frac{mass}{volume}$
	Give your answer to 2 significant figures.
	Mass of barium chloride = g [3]

Turn over © OCR 2023

17 The table shows information about three different polymers, A, B and C.

Tensile strength is the amount of load a material can take before it breaks.

	Melting point (°C)	Softening temperature (°C)	Stiffness (MPa)	Tensile strength (MPa)
Α	130	72	980	15
В	240	95	3200	65
С	250	75	2400	50

(a) A company wants to use a polymer to make a disposable cup for hot drinks.

Suggest and explain which polymer, A, B or C, the company should use.

Polymer	
Reason	
	[3]

(b) This is the repeating unit in polymer B.

$$\begin{array}{c|c} C_6H_5 & H \\ \hline \\ C & C \\ \hline \\ H & H \\ \hline \\ \\ n \end{array}$$

Draw the structure of the monomer from which polymer  ${\bf B}$  is made.

(c) Some substances are naturally occurring polymers.


Draw lines to connect each **monomer** with its naturally occurring **polymer**.

Monomer	Polymer
amino acids	DNA
nucleotides	proteins
sugars	starch

[2]

**18** A teacher investigates the reaction between hydrochloric acid and magnesium.

The diagram shows the teacher's experiment.




The teacher uses five different concentrations of hydrochloric acid. Each time they react the hydrochloric acid with 0.1 g of magnesium powder.

The table shows the teacher's results.

Concentration of hydrochloric acid (mol/dm³)	Time for magnesium powder to react (s)
0.5	117
1.0	82
1.5	48
2.0	24
2.5	16

(a)	Write the <b>balanced symbol</b> equation for the reaction between hydrochloric acid, HC <i>l</i> , and magnesium.	d
		[2]
(b)	The teacher says, 'The reaction is faster the more concentrated the hydrochloric acid'.	
	Use the results to explain why the teacher is <b>correct</b> .	
		[1]

(c) The graph shows the teacher's results.



(i) Use the graph to deduce the time for magnesium powder to react if 1.3 mol/dm<sup>3</sup> hydrochloric acid is used.

Time = .....s [1]

(ii) The teacher repeats the experiment with 0.1 g of magnesium **ribbon**.

Draw a line on the graph to show the results you would expect the teacher to get. [2]

(d)	The teacher used 0.1 g of magnesium.		
	0.1 g of magnesium reacts with hydrochloric acid to make 0.008 g of hydrogen gas.		
	Calculate the volume occupied by 0.008g of hydrogen gas in cm <sup>3</sup> .		

Relative atomic mass  $(A_r)$ : H = 1.0

Volume of hydrogen gas = ......cm<sup>3</sup> [4]

**19\*** A life-cycle assessment looks at the potential environmental impact at each stage of the life of a product.

A car manufacturer does a life-cycle assessment for cars made from

- steel
- aluminium.

The table gives information about the life-cycle impact of cars made from steel and aluminium.

	Steel	Aluminium
Production CO <sub>2</sub> emissions from mining the ore, extracting the metal, to manufacturing the car	6444 kg	9794 kg
<b>Driving</b> CO <sub>2</sub> emissions from the use of petrol or diesel	37 054 kg	36 248 kg
End of life CO <sub>2</sub> emissions saved by recycling metals rather than extracting new metals	–1546 kg	-3634 kg

Use the information in the table, and your own knowledge of how metals are extracted.

20	Butane,	C <sub>4</sub> H <sub>10</sub> ,	is	an	alkane.
----	---------	----------------------------------	----	----	---------

Butane undergoes complete combustion in oxygen.

(a)	Write the	balanced s	ymbol ed	uation for	the com	plete	combustion	of butane
-----	-----------	------------	----------	------------	---------	-------	------------	-----------

.....[2]

(b) Table 20.1 lists the energy given out when 1 g of different alkanes burn.

**Table 20.1** 

Alkane	Energy given out (kJ)
butane	49.2
ethane	52.6
methane	55.6
propane	50.4

State the relationship	between the nu	mber of carbo	n atoms in the	alkane and the	energy
given out.					

(41
 וניו

(c) Butane is found in the LPG fraction when crude oil is separated into different fractions by fractional distillation.

**Table 20.2** shows some information about four other molecules that are found in four different fractions.

**Table 20.2** 

Molecule	Formula	Boiling point (°C)
heptadecane	C <sub>17</sub> H <sub>36</sub>	302
eicosane	C <sub>20</sub> H <sub>42</sub>	342
tetracosane	C <sub>24</sub> H <sub>50</sub>	
octacosane	C <sub>28</sub> H <sub>58</sub>	436

(i) Predict the boiling point of te	rracosane	€.
-------------------------------------	-----------	----

Boiling point of tetracosane = .....°C [1]

	(ii)	Octacosane is sep molecules in <b>Table</b>	parated lower down the free <b>20.2</b> .	actionating column than	the other three
		Explain why using	ideas about intermolecu	lar forces.	
(d)			large molecules produce		-
	The	equation shows th	e cracking of octacosane	to make dodecane and	one other product.
	Con	nplete and balance	the equation.		
(e)	obta		acosane dodecane percentage supply and do		[2] lifferent fractions
		Fraction	Percentage supply (%)	Percentage demand (%)	
		LPG	2	4	
		petrol	5	23	
		naphtha	8	5	
		kerosene	12	7	
		diesel oil	17	23	
		fuel oil	56	38	
	Sug	gest and explain w	ained by cracking anothe	o obtain petrol.	
	Rea	son			

21	Met	thand	ol, CH <sub>3</sub> OH, is made in industry by reacting carbon monoxide with hydrogen.	
	This	s is th	ne equation for the reaction.	
	СО	(g) +	$2H_2(g) \rightleftharpoons CH_3OH(g)$	
	The	forw	ard reaction is exothermic.	
	(a)	A te	mperature of 250 °C and a pressure of 100 atmospheres is used for the reaction.	
		(i)	Describe and explain the effect on the yield of methanol from using a pressure of 15 atmospheres.	
		(II)		
		(ii)	The yield of methanol is greater when a temperature of 150 °C is used instead of 250 °C.	
			Suggest why a temperature of 150 °C is <b>not</b> used in industry.	
	(b)	A ca	atalyst is used to speed up the reaction.	
		Des	cribe the effect on the position of equilibrium from using a catalyst.	
				[1]
	(c)	A fa	ctory makes some methanol.	
		The of 7	y predict they will make 60 tonnes of methanol. The reaction has a percentage yield 8%.	
		Calo	culate the mass of methanol they actually make.	
			Mass of methanol = tonnes	[3]

(d) (i) Draw the structural formula of methanol,  ${\rm CH_3OH.}$ 

	(ii)	State the <b>functional group</b> in methanol.	[2]
			[1]
(e)	Met	chanol can be oxidised to methanoic acid.	
	Stat	te the <b>oxidising agent</b> used in this reaction.	
			[1]
(f)	Met	thanoic acid is a carboxylic acid.	
	Car	boxylic acids react with alcohols to form an ester and one other product.	
	(i)	What type of reaction is this?	
		Tick (✓) one box.	
		Addition	
		Condensation	
		Decomposition	
		Neutralisation	[1]
	(ii)	State the name of the other product in this type of reaction.	
			[1]

22 Group 7 elements (halogens) react with halides in solution.

Chlorine reacts with potassium bromide to form potassium chloride and bromine.

$$Cl_2(g) + 2KBr(aq) \rightarrow 2KCl(aq) + Br_2(aq)$$

(a) Describe what you would **observe** in the reaction.

[1]
-----

**(b)** This is the half equation that shows what happens to chlorine.

$$Cl_2 + 2e^- \rightarrow 2Cl^-$$

(i) Explain why this half equation shows reduction.

[1
----

(ii) Write the balanced half equation for the reaction of the bromide ions.


101

**(c)** Chlorine displaces bromine from potassium bromide because chlorine is more reactive than bromine.

Explain why chlorine is more reactive than bromine.

		[2]

(d) The diagram shows the mass spectrum for a compound containing chlorine.




State the relative molecular mass of this compound.

.....[1]

	[2
	2
	1
	State <b>two</b> advantages of instrumental methods of analysis.
(e)	chemical methods of analysis, such as mass spectrometry, have advantages over simple chemical methods of analysis.

## 23 A student plans a titration experiment.

The diagram shows some of the apparatus they use.



(a) At the end of the titration, the student reads the final volume reading from the top of the meniscus instead of from the bottom of the meniscus.

How does the measured volume of potassium hydroxide compare to the actual volume?

Tick (✓) one box.

The measured volume will be greater than the actual volume.	
The measured volume will be smaller than the actual volume.	
The measured volume will be the same as the actual volume.	

[1]

**(b)** The student uses a potassium hydroxide solution with a concentration of 0.100 mol/dm³ to neutralise the 25.0 cm³ of dilute sulfuric acid.

The table shows the student's results.

Titration number	1	2	3	4
Final burette reading (cm <sup>3</sup> )	24.1	26.6	26.0	26.8
Initial burette reading (cm <sup>3</sup> )	0.0	1.5	2.1	2.8
Titre (volume of potassium hydroxide solution used) (cm³)	24.1	25.1	23.9	24.0

(i) Calculate the average titre using the student's concordant results.

Average titre =	 $cm^3$	[1]
/ wordgo allo	 OIII	г.л

(ii) This is the equation for the reaction in this experiment.

$${\rm 2KOH} \, + \, {\rm H_2SO_4} \, \rightarrow \, {\rm K_2SO_4} \, + \, {\rm 2H_2O}$$

Calculate the concentration of sulfuric acid in g/dm<sup>3</sup>.

Relative atomic mass ( $A_r$ ): H = 1.0 O = 16.0 S = 32.0

Concentration of sulfuric acid = ......g/dm<sup>3</sup> [5]

## 26

# **ADDITIONAL ANSWER SPACE**

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).				


,		 	 
	,	 	 



#### **Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

 ${\sf OCR} \ is \ part \ of \ Cambridge \ University \ Press \ \& \ Assessment, \ which \ is \ itself \ a \ department \ of \ the \ University \ of \ Cambridge.$