



Oxford Cambridge and RSA

**For issue on or after: 13 March 2024**

**A Level Physics B (Advancing Physics)**

**H557/02 Scientific literacy in physics**

**Advance Notice Article**

**To prepare candidates for the examination taken on  
Thursday 6 June 2024 – Morning**



### INSTRUCTIONS

- Before the exam, read this article carefully and study the content of the learning outcomes for A Level Physics B (Advancing Physics).
- You can ask your teacher for advice and discuss this article with others in your class.
- You can investigate the topic of this article yourself using any resources available to you.
- Do **not** take this copy of the article or any notes into the exam.

### INFORMATION

- In the exam you will answer questions on this article. The questions are worth 20–25 marks.
- A clean copy of this article will be given to you with the question paper.
- This document has **8** pages.

### ADVICE


- In the exam you won't have time to read this article in full but you should refer to it in your answers.

## Seeing beyond the visible

- 5 Although undetectable to the human eye, objects radiate in many wavelengths beyond the visible spectrum. To better understand the nature of astronomical objects it is necessary to detect all wavelengths of electromagnetic radiation they emit. Since the middle of the twentieth century, scientists and engineers have been designing telescopes to detect these non-visible wavelengths.

The atmosphere is transparent to visible light and radio waves as can be seen in **Fig. 1**. For the purposes of this article, the height of the atmosphere can be taken to be 140 km.

**Fig. 1**




- 10 Observations can be made with ground-based telescopes in the visible and radio wavelengths, but to observe in other bands of the spectrum requires telescopes either to be as high up as possible, to minimise the effects of the atmosphere, or launched into space to completely escape the atmosphere.

## A problem of resolution

- 15 The angular resolution of an optical instrument is the minimum angle that can be resolved due to the diameter of the collecting lens (in binoculars, say) or mirror (in large optical and infrared telescopes and dish radio telescopes) as shown in **Fig. 2**.

**Fig. 2**



The angular resolution in radians is given by the equation  $angular\ resolution = \frac{1.2\lambda}{d}$  where  $\lambda$  is  
20 the wavelength of the radiation and  $d$  is the diameter of the collecting lens or mirror.

The dependence of resolution on wavelength means that radio receivers, such as the Jodrell Bank telescope, need very large diameter mirrors. The Jodrell Bank dish shown in **Fig. 3** has a diameter of 76.2 m. It collects radio signals of wavelength 0.06 m. The minimum angular separation it can resolve is about four times that of the unaided eye.

25 **Fig. 3** Jodrell Bank main dish




### Telescopes in space

The Chandra X-ray Observatory was carried into space by the Space Shuttle in July 1999. It was placed in a highly elliptical orbit, varying its distance from Earth from approximately  $1.6 \times 10^7$  m (at closest approach) to  $1.5 \times 10^8$  m (at its furthest from Earth). X-rays are emitted  
30 from highly energetic objects such as matter spiralling into black holes and the Chandra has made important observations of X-rays emitted from the region around the super-massive black hole at the centre of our galaxy.


The James Webb telescope (JWST), the largest telescope yet taken into space, was launched on December 25<sup>th</sup> 2021 from Kourou, French Guiana. It is placed about  $1.5 \times 10^9$  m beyond Earth in line with Earth and the Sun as illustrated in **Fig. 4**. At this point, known as the L2 point, the gravitational forces from the Sun and Earth combine to give sufficient force on the telescope so that its orbital period around the Sun is the same as that of Earth even though it is further away. In fact, the telescope is not actually on the L2 point but orbits around it as indicated on the diagram. The radius of the orbit around the L2 point is about  $8 \times 10^6$  m.

40 **Fig. 4** (not to scale)



The JWST is an infrared telescope which detects electromagnetic radiation at wavelengths from 600 nm to  $29 \mu\text{m}$ .

**Fig. 5**




45 It is important to keep the science instruments as cool as possible. To achieve this, the telescope always faces away from Earth and the Sun. The sunshield reflects much of the radiation and cooling systems keep the operating temperature of the instruments below 50 K.

One of these instruments is a near infrared camera (NIRCam) which can image in two wavelength ranges; from 600 nm to 2300 nm (short wavelength) and 2400 nm to 5000 nm (long wavelength). The short wavelength range uses eight individual sensitive surfaces each of  $2040 \times 2040$  pixels.

- 50 In the short wavelength range, the eight surfaces can be combined to give a field of view of approximately  $1.2 \times 10^{-3} \text{ rad} \times 6.4 \times 10^{-4} \text{ rad}$ .

**Fig. 6** Field of view of eight combined sensitive surfaces (simplified)



- 55 The near infrared spectrograph (NIRSpec) covers the same range of wavelengths as NIRC and is used to produce spectra of objects of interest. Absorption lines in spectra identify the chemicals present in the source and, through redshift measurements, can give information about the distance to the source.

For objects moving away from Earth at velocities much less than  $c$ :

$$\text{fractional increase in wavelength} = \frac{\lambda_{\text{observed}} - \lambda_{\text{emitted}}}{\lambda_{\text{emitted}}} = \frac{v}{c}$$

where  $v$  is the velocity of the light source (for example, a galaxy) and  $c$  is the velocity of light.

- 60 Hubble's Law states: velocity of recession =  $H_0 d$ , where  $H_0$  is the Hubble constant and  $d$  is the distance to the receding object, hence the distance to the object can be found from its redshift.

### Seeing through the clouds

The space between stars is a near-vacuum. Interstellar space has roughly one million atoms per  $\text{m}^3$ , a factor of  $10^{19}$  less than the atmosphere of Earth.

- 65 However, there are regions of space with greater density of gas. Embedded in these gas clouds are grains of dust of diameter about 500 nm composed of carbon, ice and other chemicals.

- 70 These grains can absorb, scatter and emit electromagnetic radiation. Radiation of wavelengths greater than the size of the dust grains has little interaction with the dust so whereas short wavelength radiation is scattered by interstellar dust grains, longer wavelengths can pass through. This means that it is possible to observe behind clouds of dust with infrared radiation. This will help the telescope look deep into space and image some of the oldest galaxies, revealing the evolution of the early Universe. It will also, using NIRSpec, record the chemical compositions of planets in the Solar System and beyond. These observations could give evidence of the potential for life on other worlds.

- 75 From our astronomical 'backyard' to the furthest reaches of space, images and data from the JWST have already provided a wealth of information to keep researchers busy for years to come and will help to further our understanding of the Universe.





---

**OCR**  
Oxford Cambridge and RSA

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series. If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.