



Oxford Cambridge and RSA


Thursday 20 June 2024 – Afternoon

A Level Mathematics B (MEI)

H640/03 Pure Mathematics and Comprehension

Insert

Time allowed: 2 hours



INSTRUCTIONS

- Do **not** send this Insert for marking. Keep it in the centre or recycle it.

INFORMATION

- This Insert contains the article for Section B.
- This document has **4** pages.


Tangents and normals to a quadratic curve

Tangents

Fig. C1 shows the curve $y = x^2$ together with tangents to the curve at points A $(-3, 9)$ and B $(1, 1)$. The tangents cross at the point $(-1, -3)$. This has x -coordinate -1 , which is the mean of the x -coordinates of points A and B.

Fig. C1

5



For the curve $y = x^2$, the equation of the tangent at a general point (t, t^2) is $y = 2tx - t^2$. So the equation of the tangent at the point (t_1, t_1^2) is $y = 2t_1x - t_1^2$. There is a similar equation for the tangent at the point (t_2, t_2^2) , and these two tangents cross where $2t_1x - t_1^2 = 2t_2x - t_2^2$.

This gives $2x(t_1 - t_2) = t_1^2 - t_2^2$ so $2x(t_1 - t_2) = (t_1 - t_2)(t_1 + t_2)$ hence $x = \frac{t_1 + t_2}{2}$. The y -coordinate of the point of intersection is $t_1 t_2$.

10

The general quadratic curve has equation $y = ax^2 + bx + c$. The tangents at any two points P and Q on this curve also cross at a point whose x -coordinate is equal to the mean of the x -coordinates of P and Q. So if P has x -coordinate x_P and Q has x -coordinate x_Q then the x -coordinate of the intersection point of the tangents is $\frac{x_P + x_Q}{2}$. The y -coordinate of the intersection point can be

shown to be $ax_Px_Q + b\left(\frac{x_P+x_Q}{2}\right) + c$. This is equivalent to

$a\left(\frac{x_P+x_Q}{2}\right)^2 + b\left(\frac{x_P+x_Q}{2}\right) + c - a\left(\frac{x_P-x_Q}{2}\right)^2$. The formula $ax_Px_Q + b\left(\frac{x_P+x_Q}{2}\right) + c$ looks simpler but $y = a\left(\frac{x_P+x_Q}{2}\right)^2 + b\left(\frac{x_P+x_Q}{2}\right) + c - a\left(\frac{x_P-x_Q}{2}\right)^2$ is in terms of the x -coordinate of the point of intersection, apart from the last term. For pairs of points with $x_P - x_Q = h$ where h is a constant, the point of intersection of the tangents lies on the curve $y = ax^2 + bx + c - \frac{ah^2}{4}$. This curve is a translation of the original curve $y = ax^2 + bx + c$.


15

20

Normals

Fig. C2 shows the curve $y = x^2$ together with normals to the curve at points A $(-3, 9)$ and B $(1, 1)$. The normals cross at the point $(-12, 7.5)$.

Fig. C2



For the curve $y = x^2$, the coordinates of the point of intersection are not as simply related to the coordinates of A and B as in the case of the tangents. The equation of the normal at the point (t, t^2) is $y = -\frac{x}{2t} + t^2 + \frac{1}{2}$. The normals at points (t_1, t_1^2) and (t_2, t_2^2) cross when $x = -2t_1t_2(t_1 + t_2)$ and $y = t_1^2 + t_2^2 + t_1t_2 + \frac{1}{2}$.

25

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series. If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.