

Friday 21 June 2024 - Morning

A Level Chemistry A

H432/03 Unified chemistry

Time allowed: 1 hour 30 minutes

You must have:

· the Data Sheet for Chemistry A

You can use:

- · a scientific or graphical calculator
- an HB pencil



331429 33

29 331429

29 331429 331429 331⁴²⁹ 331429 3314 29 331429

29 331429

Please write clearly in black ink. Do not write in the barcodes.										
Centre number						Candidate number				
Circt name (a)										
First name(s)										
Last name										

29 331429

INSTRUCTIONS

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.
- Answer all the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

INFORMATION

- The total mark for this paper is **70**.
- The marks for each question are shown in brackets [].
- Quality of extended response will be assessed in questions marked with an asterisk (*).
- This document has **20** pages.

ADVICE

Read each question carefully before you start your answer.



1	This question is about acids, bases and salts.		
(a)	What is the pH of 1.00 dm ³ of 0.400 mol dm ⁻³ of NaOH(aq) at 298 K?		
		pH =	[2]
(b)	Water is added to $10.0 \mathrm{cm^3}$ of $0.750 \mathrm{moldm^{-3}}$ HC $l(\mathrm{aq})$ to produce $100 \mathrm{cm^3}$ or	f diluted HC <i>l</i> (aq).	
` ,	What is the pH of the diluted $HCl(aq)$?	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	
	Give your answer to 2 decimal places.		
		pH =	[1]
(c)	A solution has concentrations of 0.300 mol dm $^{-3}$ CH $_3$ COOH(aq) and 0.100 m CH $_3$ COONa(aq).	oldm ⁻³	
	$K_{\rm a}$ for CH ₃ COOH = 1.75 × 10 ⁻⁵ mol dm ⁻³ at 298 K.		
	What is the pH of the solution at 298 K?		
	Give your answer to 2 decimal places.		
		pH =	[21
		L	·1

(d)	A student is provided with hydrated copper(II) nitrate, Cu(NO ₃) ₂ •3H ₂ O.
	The student needs to prepare a standard solution of $Cu(NO_3)_2$ •3 H_2O with a concentration of 0.200 mol dm ⁻³ . The student has access to usual laboratory apparatus and equipment.
	Describe how the student would prepare $100.0\mathrm{cm^3}$ of this solution, giving quantities, apparatus and method.

_	T1 :	e e			11.00				
2	I nis (question	IS	about	aiπerent	areas	ΟŤ	chemistry	٧.

(a)	Hydrogen gas is manufactured by the chemical industry from the reversible reaction of methane
	and steam, shown below.

$$CH_4(g) + H_2O(g) \rightleftharpoons 3H_2(g) + CO(g)$$
 $\Delta H = +195 \text{ kJ mol}^{-1}$

Average bond enthalpies are shown in the table.

Bond	H–H	O–H	C≡O
Average bond enthalpy /kJ mol ⁻¹	+436	+464	+1077

(i)	Why do all average bond enthalpies have a positive value?							
		[1]						
(ii)	Determine the C–H bond enthalpy, in kJ mol ⁻¹ , using the information above.							

()

C-H bond enthalpy =
$$kJ mol^{-1}$$
 [3]

(iii) Hydrogen gas is being considered as a household fuel to replace methane.

The enthalpy change of formation, $\Delta_{\rm f}H,$ for $\rm H_2O(I)$ is $-285.8\,kJ\,mol^{-1}.$

Determine the energy released when $60.0\,\mathrm{m}^3$ of hydrogen is used as a household fuel at RTP.

Give your answer to $\bf 3$ significant figures and in $\bf standard\ form.$

		5
(b)	Compou	nd A is a chloride of a Period 3 element.
	A studer	at carries out the 2 steps below to find the formula of compound A .
	Step 1	The student adds 5.00×10^{-4} mol of compound A to water. A colourless solution is formed.
	Step 2	The colourless solution reacts with exactly $60.0\mathrm{cm^3}$ of $2.50\times10^{-2}\mathrm{moldm^{-3}}$ AgNO $_3(\mathrm{aq})$ to form a white precipitate.
(i)	Write an	ionic equation, with state symbols, for the reaction in Step 2 .
		[1]
(ii)	Determin	ne the formula of compound A .
		formula of A =[3]

Turn over © OCR 2024

(c) Compound B, shown below, is an antiviral medicine.

compound B

(i)	What is the molecular formula of compound B ?

(ii) How many chiral carbon atoms are there in one molecule of compound **B**?

......[1]

.....[1]

- (iii) A research chemist synthesises two related compounds, compound **C** and compound **D**, from compound **B**.
 - In compound **C**, the N atoms in compound **B** had been replaced by P atoms.
 - In compound **D**, the O atoms in compound **B** had been replaced by S atoms.

What is the difference between the relative molecular masses of compound C and compound D?


difference =[2]

7 BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

Turn over for the next question

- This question is about compounds and ions of iron(II) and iron(III) that contain ethanedioate ions, $C_2O_4^{\ 2-}$.
- (a) The ${\rm C_2O_4^{\,2-}}$ ion, shown below, is an example of a bidentate ligand.



(i)	Explain what is meant by the term bidentate ligand .						
	[2]						

(ii) A complex ion ${\bf E}$ contains three ${\rm C_2O_4^{2-}}$ ions bonded to an iron(III) ion in an octahedral shape.

Complex ion **E** exists as a mixture of two optical isomers.

Draw 3D diagrams to show the structures of the optical isomers of **E**.

Include any overall charge.

(b)	A student plans an investigation to find the number of waters of crystallisation, \mathbf{x} , in a sample of hydrated iron(II) ethanedioate, $\text{FeC}_2\text{O}_4 \bullet \mathbf{x}\text{H}_2\text{O}$.	
	The student decides to carry out a redox titration between solutions of iron(II) ethanedioate and potassium manganate(VII) in acidic conditions.	
(i)	In the titration, both iron(II) ions and ethanedioate, $\rm C_2\rm O_4^{2-}$, ions are oxidised.	
	Construct half-equations for the oxidation of iron(II) and ethanedioate ions.	
	Oxidation of iron(II) ions	
	Oxidation of ethanedioate ions	
	ŗ	 21
	L ⁱ	2]

(ii) The student prepares a 250.0 cm³ solution of iron(II) ethanedioate by dissolving 1.295 g of FeC₂O₄•xH₂O, in dilute sulfuric acid.

The student titrates $25.0\,\mathrm{cm^3}$ samples of this solution with $0.0200\,\mathrm{mol\,dm^{-3}}$ KMnO₄ in the burette. The student carries out a trial, followed by three further titrations.

The diagrams show the initial burette readings and the final burette readings for the student's three further titrations.

Titrat	tion 1	Titrat	tion 2	Titration 3			
Initial reading	Final reading	Initial reading	Final reading	Initial reading	Final reading		
	=	======================================	= 45 = 46 = 47 = 47		======================================		

All burette readings are measured to the nearest 0.05 cm³.

Complete the titration table.

	1	2	3
Final reading/cm ³			
Initial reading/cm ³			
Titre/cm ³			

[3]

(iii) The uncertainty in each burette reading is $\pm 0.05 \, \text{cm}^3$.

Calculate the percentage uncertainty for the titre in **Titration 1**.

percentage uncertainty = % [1]

© OCR 2024

(iv)* In the titration, 5 mol of iron(II) ethanedioate reacts with 3 mol of manganate(VII) ions. Analyse the student's results to find the number of waters of crystallisation, x, in the hydrated iron(II) ethanedioate, FeC₂O₄•**x**H₂O. [6] Extra answer space if required.

.....

4	This question is about the chemistry of compounds containing phosphorus.	
(a)	Phosphorus forms several acids including H ₃ PO ₄ and H ₃ PO ₃ .	
	H ₃ PO ₄ is a tribasic acid. The equilibria for the dissociations are shown below.	
	1 $H_3PO_4 \rightleftharpoons H^+ + H_2PO_4^-$	
	2 $H_2PO_4^- \rightleftharpoons H^+ + HPO_4^{2-}$	
	3 $HPO_4^{2-} \rightleftharpoons H^+ + PO_4^{3-}$	
(i)	During the equilibria, H ₂ PO ₄ ⁻ behaves both as an acid and as a base.	
	Explain this statement, using the equilibria 1, 2 and 3, as required.	
		[2]
(ii)	In a $\rm H_3PO_3$ molecule, the O atoms are covalently bonded to the P atom. The H atoms are bonded to the O atoms.	
	Draw the structure of a H ₃ PO ₃ molecule, showing all the bonds.	
	On your diagram, add the values for the O–P–O and P–O–H bond angles.	
		101
,,,, ,		[3]
(iii)	The systematic name of H ₃ PO ₄ is phosphoric(V) acid.	

.....[1]

What is the systematic name of H_3PO_3 ?

(b)	Phosphine, PH ₃ , is a poisonous gas.	
(i)	Phosphine reacts with oxygen gas to form phosphorus(V) oxide and water.	
	Write the equation for this reaction.	
		. [1]
(ii)	Aqueous silver nitrate, AgNO ₃ , is reduced by PH ₃ . The unbalanced equation is shown below.	
	Balance the equation and use oxidation numbers to explain why this is a redox reaction.	
	Explanation	
		[31

(c)	When phosphorus(V) chloride, PCl_5 , and ammonium chloride are heated together, the compound $P_3N_3Cl_6$ is formed, together with HCl gas.
	${\rm P_3N_3C}l_6$ has a cyclic structure, like the Kekulé structure of benzene.
(i)	Write an equation for the reaction of $\mbox{PC}l_5$ and ammonium chloride to form $\mbox{P}_3\mbox{N}_3\mbox{C}l_6$.
	[1]
(ii)	Calculate the percentage by mass of P in $P_3N_3Cl_6$.
	Give your answer to 2 decimal places.
	percentage by mass of P = % [2]
(iii)	Suggest one example of evidence that could show that $P_3N_3Cl_6$ has a Kekulé structure rather than a delocalised structure.
	[1]
(iv)	In a molecule of $\rm P_3N_3C\mathit{l}_6$ all the N and C l atoms are bonded to P atoms.
	Suggest a possible structure for a molecule of $P_3N_3Cl_6$.

15 BLANK PAGE


PLEASE DO NOT WRITE ON THIS PAGE

Turn over for the next question

- **5** This question is about the analysis of organic compounds.
- (a) A student investigates the alkaline hydrolysis of 1-bromopropane as outlined below.
 - **Step 1** The student adds 1-bromopropane to an excess of aqueous potassium hydroxide, KOH(aq), in a pear-shaped flask.
 - **Step 2** A TLC chromatogram is run using propan-1-ol and the reaction mixture.
 - **Step 3** The reaction mixture is refluxed.

 A TLC chromatogram of the reaction mixture is run every 10 minutes.

The TLC chromatograms are shown below.



(i)	Determine the $R_{\rm f}$ value of propan-1-ol.
	Show your working.
	R _f =[1]
(ii)	Write an equation for the alkaline hydrolysis of 1-bromopropane.
	Show structures of organic compounds.
	[1]
(iii)	A student investigates the alkaline hydrolysis of 1-chloropropane using the same method as for 1-bromopropane.
	Predict, with reasons, how the appearance of the reaction mixture in the chromatogram produced after 20 minutes would be different when 1-chloropropane is used instead of 1-bromopropane.
	Suggest why propan-1-ol is run alongside the reaction mixture.
	[3]
	[5]

(b) Compounds F, G, H and I are structural isomers.

A student carries out test-tube tests on the compounds. The student records the observations after carrying out each test. These are shown in **Table 5.1.**

In **Table 5.1**, 2,4-dintrophenylhydrazine has been abbreviated to 2,4-DNP.

Table 5.1

	Test			
Compound	2,4-DNP	Acidified dichromate(VI) reflux	Bromine water	Tollens' reagent
F	Orange solution	Green solution	Colourless solution	Colourless solution
G	Orange solution	Green solution	Orange solution	Colourless solution
Н	Orange precipitate	Orange solution	Orange solution	Colourless solution
I	Orange precipitate	Green solution	Orange solution	Silver mirror

(i)	Write the formula of the species causing the colours after refluxing with acidified dichromate(VI).
	Green solution
	Orange solution
	[2]

- (ii)* The student is provided with further information about compounds F-I.
 - They all have the molecular formula ${\rm C_5H_{10}O}.$ One of the compounds is alicyclic.

 - The other compounds are unbranched.

Use this further information and the student's observations in **Table 5.1** to answer the following.

- How do the observations provide evidence for the possible functional groups in compounds **F-I**?
- Suggest a possible structure for each of the compounds **F–I**.

Show your reasoning.	[6]
Extra answer space if required.	

EXTRA ANSWER SPACE

If you need extra space use this lined page. You must write the question numbers clearly in the margin.		
•••••		
•••••		



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.