



Oxford Cambridge and RSA

Wednesday 18 June 2025 – Afternoon

A Level Further Mathematics B (MEI)

Y434/01 Numerical Methods

Time allowed: 1 hour 15 minutes



You must have:

- the Printed Answer Booklet
- the Formulae Booklet for Further Mathematics B (MEI)
- a scientific or graphical calculator

QP

INSTRUCTIONS

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined page at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION

- The total mark for this paper is **60**.
- The marks for each question are shown in brackets [].
- This document has **12** pages.

ADVICE

- Read each question carefully before you start your answer.

- 1 The table shows some values of x and the associated values of $f(x)$.

x	1.9	2.0	2.1
$f(x)$	0.216 92	0.2	0.184 84

(a) Determine an estimate of $f'(2.0)$ using the forward difference method. [2]

(b) Determine an estimate of $f'(2.0)$ using the central difference method. [2]

- 2 The numbers p and q are approximated by

$$P = 323 \text{ and } Q = 162.$$

P has been found by **rounding** p to the nearest whole number.

Q has been found by **chopping** q to the nearest whole number.

(a) (i) Find the maximum possible relative error in using P to approximate p . [1]

(ii) Find the maximum possible relative error in using Q to approximate q . [1]

(b) Determine the range of possible values of $R = \frac{200}{p-2q}$. [3]

(c) Explain why your answer to part (b) is so large. [1]

- 3 Approximations to $\int_{0.5}^{1.3} \sqrt{1+x^3} \, dx$ using the midpoint rule, the trapezium rule and Simpson's rule with $n = 1$ and $n = 2$ are shown in the table. The table is incomplete.


n	M_n	T_n	S_{2n}
1			1.081 111
2	1.074 256		

(a) Complete the copy of the table in the **Printed Answer Booklet**. Give your answers to 6 decimal places. [4]

(b) **Without doing any further calculations**, state the value of $\int_{0.5}^{1.3} \sqrt{1+x^3} \, dx$ as accurately as possible. You must justify the precision quoted. [1]


- 4 The Newton-Raphson method is to be used to find the positive root of the equation $\tanh x - x^2 + 4 = 0$.

The diagram shows part of the graph of $y = \tanh x - x^2 + 4$.



- (a) On the copy of the diagram in the **Printed Answer Booklet**, show how the Newton-Raphson method works to find x_1 using the starting value $x_0 = 1$. [1]
- (b) Use the Newton-Raphson method using the starting value $x_0 = 1$ to determine the values of x_1 and x_2 correct to **8** decimal places. [4]
- (c) Continue the iteration to determine the value of the positive root of the equation $\tanh x - x^2 + 4 = 0$ correct to **7** decimal places. [2]

- 5 The diagram shows part of the graph of $y = \sinh x - 3x + 1$.



The largest positive root, α , of the equation $\sinh x - 3x + 1 = 0$ is to be found using the secant method with $x_0 = 2$ and $x_1 = 3$.

Table 5.1 shows the associated spreadsheet output.

Table 5.1

	A	B	C	D	E
2	r	x_r	$f(x_r)$	x_{r+1}	$f(x_{r+1})$
3	0	2	-1.37314	3	2.01787
4	1	3	2.017875	2.404935	-0.7211
5	2	2.404935	-0.72109	2.561598	-0.2451
6	3	2.561598	-0.24513	2.642285	0.06018
7	4	2.642285	0.060175	2.626382	-0.0035
8	5	2.626382	-0.00348	2.627252	-5E-05
9	6	2.627252	-4.5E-05	2.627264	3.5E-08

- (a) Write down a suitable formula for cell D4. [2]
- (b) State the most accurate approximation to α in the spreadsheet output. [1]
- (c) Determine whether this approximation is accurate to 6 decimal places. [2]
- (d) Explain what would happen if the initial entry in cell D3 were to be changed from 3 to 1. [1]

Table 5.2 shows some further analysis of the sequence of approximations to α .

Table 5.2

A	B	C	D	E
r	x_r	$x_{r+1} - x_r$	$\frac{x_{r+2} - x_{r+1}}{x_{r+1} - x_r}$	$\frac{x_{r+2} - x_{r+1}}{(x_{r+1} - x_r)^2}$
1	3	-0.5951	-0.2633	0.44242
2	2.404935	0.15666	0.51504	3.28756
3	2.561598	0.08069	-0.1971	-2.4427
4	2.642285	-0.0159	-0.0547	3.44193
5	2.626382	0.00087	0.01315	15.1113
6	2.627252	1.1E-05		
7	2.627264			

(e) Explain what may be inferred from the entries in column D about the order of convergence of this sequence of approximations to α . [2]

(f) Explain what may be inferred from the entries in column E about the order of convergence of this sequence of approximations to α . [2]

6 The midpoint rule is used to find a sequence of approximations to $\int_0^1 \sqrt{\sin x} dx$.

The associated spreadsheet output, together with some further analysis, is shown in the table.

n	M_n	difference	ratio
1	0.6924056		
2	0.6615057	-0.0308999	
4	0.6498274	-0.0116782	0.37794
8	0.6454773	-0.0043501	0.3725
16	0.6438811	-0.0015962	0.36692
32	0.643302	-0.0005791	0.36281
64	0.6430936	-0.0002085	0.35996
128	0.6430189	-7.463E-05	0.35801

(a) Explain what the entries in the ratio column tell you about the order of the midpoint rule in this case. [2]

(b) Use the information in the table to determine the most accurate estimate of $\int_0^1 \sqrt{\sin x} dx$ possible. You must justify the precision quoted. [5]

- 7 The equation $8 \ln(x+2) - 3x + 1 = 0$ has a positive root, γ , and a negative root, δ .

It is proposed that the iterative formula

$$x_{n+1} = g(x_n), \text{ where } g(x) = 8 \ln(x+2) - 2x + 1,$$

should be used to find γ .

It is found that $g'(\gamma) \approx -0.977$.

- (a) Explain what this tells you about the speed of convergence of the sequence of approximations to γ which would be generated by $x_{n+1} = g(x_n)$. [2]

The relaxed iteration

$$x_{n+1} = (1 - \lambda)x_n + \lambda g(x_n) \text{ with } \lambda = 0.506$$

is used to find a sequence of approximations to γ .

The spreadsheet output is shown in the table, together with the values of $f(x_r) = 8 \ln(x_r + 2) - 3x_r + 1$ for $r = 0, 1, 2, 3, 4$.

	E	F	G	
1	r	x_r	$f(x_r)$	
2	0	5	1.567281192	
3	1	5.7930443	0.046719768	
4	2	5.8166845	3.04288E-05	
5	3	5.8166999	-4.07401E-09	
6	4	5.8166999	5.47118E-13	

- (b) State the value of γ as accurately as you can, justifying the precision quoted. [1]
- (c) Explain why the values displayed in cells F5 and F6 are the same as each other, but the values in cells G5 and G6 are different to each other. [2]

The negative root of the equation $8 \ln(x+2) - 3x + 1 = 0$, δ , is close to -1 .

It is found that $g'(\delta) \approx 13.9$.

- (d) Explain why the iterative formula

$$x_{n+1} = g(x_n) \text{ with } x_0 = -1$$

may **not** be used to find δ . [1]

(e) (i) Use the relaxed iteration

$$x_{n+1} = (1 - \lambda)x_n + \lambda g(x_n) \text{ with } \lambda = 0.506 \text{ and } x_0 = -1$$

to find x_1, x_2, x_3 and x_4 , giving your answers correct to **3** decimal places. **[2]**

(ii) State what would happen if the iteration in part **(e)(i)** were to be continued. **[1]**

(f) Use the relaxed iteration

$$x_{n+1} = (1 - \lambda)x_n + \lambda g(x_n) \text{ with } \lambda = -0.078 \text{ and } x_0 = -1$$

to determine the value of δ correct to **7** decimal places. **[2]**

- 8 Following an injury an athlete returns to training in preparation for a regional competition. At the end of each week the athlete's trainer records the time, t seconds, the athlete takes to run 100 metres.

Table 8.1 shows some of the times recorded at the end of week W .

Table 8.1

W	0	2	3
t	18.00	17.68	16.83

The trainer uses the information in the table to construct a polynomial of degree 2 to model these data.

- (a) Use Lagrange's interpolating polynomial of degree 2 to obtain the trainer's model. [3]

Subsequently it is found that when $W = 4$, $t = 15.44$.

- (b) Determine whether the model is a good fit for these values. [1]

The trainer decides to use the three most recent data points, when $W = 2, 3$ and 4, to construct a new model using Newton's forward difference interpolation formula.

Table 8.2 shows the difference table for these data.

Table 8.2

W	t		
2	17.68		
		-0.85	
3	16.83		-0.54
		-1.39	
4	15.44		

- (c) The trainer obtains the interpolation formula

$$t = -0.27W^2 + BW + C, \text{ where } B \text{ and } C \text{ are constants.}$$

- Determine the value of B .
- Determine the value of C . [3]

- (d) Explain why the new model is a refinement of the model found in part (a). [1]

At the regional competition, the athlete needs to run 100 metres in less than 11 seconds in order to qualify for a national competition. The regional competition takes place when $W = 6$.

- (e) Determine whether, according to the new model, the athlete will qualify for the national competition. [1]
- (f) Explain whether it would be appropriate to use the new model to predict the athlete's times for $W \geq 7$. [1]

END OF QUESTION PAPER

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series. If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.