

OXFORD CAMBRIDGE AND RSA EXAMINATIONS

Wednesday 4 June 2025 – Afternoon

A Level Mathematics A

H240/01 Pure Mathematics

Time allowed: 2 hours

plus your additional time allowance

YOU MUST HAVE:

**the Printed Answer Booklet or any suitable paper
provided by the centre. The Printed Answer Booklet may
be enlarged by the centre.**

a scientific or graphical calculator

READ INSTRUCTIONS OVERLEAF



INSTRUCTIONS

Use black ink. You can use an HB pencil, but only for graphs and diagrams.

If you use the Printed Answer Booklet write your answer to each question in the space provided in the PRINTED ANSWER BOOKLET. If you need extra space use the lined page at the end of the Printed Answer Booklet. The question numbers must be clearly shown.

If you use the Printed Answer Booklet fill in the boxes on the front of the Printed Answer Booklet.

Answer ALL the questions.

Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question.

The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. When a numerical value is needed use $g = 9.8$ unless a different value is specified in the question.

Do NOT send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION

The total mark for this paper is 100.

The marks for each question are shown in brackets [].

ADVICE

Read each question carefully before you start your answer.

FORMULAE

A Level Mathematics A (H240)

Arithmetic series

$$S_n = \frac{1}{2}n(a + l) = \frac{1}{2}n\{2a + (n - 1)d\}$$

Geometric series

$$S_n = \frac{a(1 - r^n)}{1 - r}$$

$$S_\infty = \frac{a}{1 - r} \quad \text{for } |r| < 1$$

Binomial series

$$(a + b)^n = a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_r a^{n-r}b^r + \dots + b^n$$

$(n \in \mathbb{N})$

where ${}^nC_r = {}_nC_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$

$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots$$

$$(|x| < 1, n \in \mathbb{R})$$

Differentiation

$f(x)$	$f'(x)$
$\tan kx$	$k \sec^2 kx$
$\sec x$	$\sec x \tan x$
$\cot x$	$-\operatorname{cosec}^2 x$
$\operatorname{cosec} x$	$-\operatorname{cosec} x \cot x$

Quotient rule $y = \frac{u}{v}, \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$

Differentiation from first principles

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Integration

$$\int \frac{f'(x)}{f(x)} dx = \ln|f(x)| + c$$

$$\int f'(x) (f(x))^n dx = \frac{1}{n+1} (f(x))^{n+1} + c$$

Integration by parts $\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$

Small angle approximations

$\sin \theta \approx \theta$, $\cos \theta \approx 1 - \frac{1}{2}\theta^2$, $\tan \theta \approx \theta$ where θ is measured in radians

Trigonometric identities

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \quad \left(A \pm B \neq \left(k + \frac{1}{2}\right)\pi \right)$$

Numerical methods

Trapezium rule:

$$\int_a^b y \, dx \approx \frac{1}{2}h \{(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1})\},$$

$$\text{where } h = \frac{b - a}{n}$$

The Newton-Raphson iteration for solving $f(x) = 0$:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

Probability

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cap B) = P(A)P(B | A) = P(B)P(A | B)$$

$$\text{OR } P(A | B) = \frac{P(A \cap B)}{P(B)}$$

Standard deviation

$$\sqrt{\frac{\sum (x - \bar{x})^2}{n}} = \sqrt{\frac{\sum x^2}{n} - \bar{x}^2} \quad \text{OR} \quad \sqrt{\frac{\sum f(x - \bar{x})^2}{\sum f}} = \sqrt{\frac{\sum fx^2}{\sum f} - \bar{x}^2}$$

The binomial distribution

If $X \sim B(n, p)$ then $P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x}$,

mean of X is np , variance of X is $np(1 - p)$

Hypothesis test for the mean of a normal distribution

If $X \sim N(\mu, \sigma^2)$ then $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$ and $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$

Percentage points of the normal distribution

If Z has a normal distribution with mean 0 and variance 1 then, for each value of p , the table gives the value of z such that $P(Z \leq z) = p$.

p	0.75	0.90	0.95	0.975	0.99	0.995
z	0.674	1.282	1.645	1.960	2.326	2.576

p	0.9975	0.999	0.9995
z	2.807	3.090	3.291

Kinematics

Motion in a straight line

$$v = u + at$$

$$s = ut + \frac{1}{2}at^2$$

$$s = \frac{1}{2}(u + v)t$$

$$v^2 = u^2 + 2as$$

$$s = vt - \frac{1}{2}at^2$$

Motion in two dimensions

$$\mathbf{v} = \mathbf{u} + \mathbf{a}t$$

$$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$$

$$\mathbf{s} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$$

$$\mathbf{s} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2$$


1 Express each of the following in the form px^q , where p and q are constants.

(a) $\frac{2}{\sqrt[4]{x}}$ [1]

(b) $(5x\sqrt{x})^3$ [2]

(c) $\sqrt{2x^3} \times \sqrt{8x^5}$ [2]

(d) $x^5(27x^6)^{\frac{1}{3}}$ [3]



The diagram shows the curve $y = ax(x - b)$, where a and b are constants and $b > 0$.

- (a) Given that the curve has a stationary point at $x = 3$, state the value of b . [1]
- (b) Given also that the stationary point at $x = 3$ is a maximum, state what can be deduced about the value of a . [1]
- (c) Find the y -coordinate of the stationary point, giving your answer in terms of a . [1]
- (d) State the range of values of x for which the curve is increasing. [1]

- 3 (a) A sequence has terms $u_1, u_2, u_3, \dots$ defined by $u_1 = 3$ and $u_{n+1} = u_n^2 - 5$ for $n \geq 1$.
- (i) Find the values of u_2, u_3 and u_4 . [2]
- (ii) Describe the behaviour of the sequence. [1]
- (b) The second, third and fourth terms of a geometric progression are 12, 8 and $\frac{16}{3}$.
- Determine the sum to infinity of this geometric progression. [3]
- 4 The position vector of the point A is $0.5\mathbf{i} - 0.5\mathbf{j}$.
- (a) Determine whether $0.5\mathbf{i} - 0.5\mathbf{j}$ is a unit vector. [2]
- (b) Find the direction of $0.5\mathbf{i} - 0.5\mathbf{j}$, giving your answer in degrees. [2]

The position vector of the point B is $3.5\mathbf{i} + c\mathbf{j}$, where c is a constant.

- (c) Given that $|\overrightarrow{AB}| = 5$, determine the possible values of c . [3]

- 5 Scientists are comparing h , the average height of a child in cm, with t , the age of the child in years. They suggest the model $h = at + b$ for $t \geq 2$, where a and b are constants.**

They find that the average height of a 2 year old child is 87 cm and the average height of a 5 year old child is 108 cm.

- (a) Find the values of a and b that are consistent with the scientists' findings. [4]**


- (b) (i) Sam is 4 years old.**

Use the model to predict Sam's height. [2]

- (ii) Comment on the accuracy of this prediction. [1]**

- (c) Suggest ONE possible limitation of this model when predicting the average height of a 12 year old child. [1]**

NOT TO SCALE



The diagram shows a triangle ABC . The arc BD is part of a circle with centre A and radius 6 cm .

The area of the sector ABD is 14.4 cm^2 .

(a) Show that angle BAD is 0.8 radians. [1]

The area of the triangle ABC is THREE times the area of the sector ABD .

(b) Find the length AC . [2]

(c) Find the perimeter of the region BCD . [5]

7 In this question you must show detailed reasoning.

**A curve has parametric equations $x = t^3 + t^2$,
 $y = t^2 + 2t$ for all real values of t .**

**The curve passes through the point P with
coordinates $(2, 3)$.**

**(a) Show that the equation of the tangent to the curve
at the point P can be written as $5y = 4x + 7$. [5]**

**(b) Determine the coordinates of the point where the
tangent to the curve at P meets the curve again. [4]**

**8 (a) Find the first THREE terms in the expansion of
 $(2 + x)^{-3}$ in ascending powers of x . [4]**

**(b) Hence find the first THREE terms in the expansion
of $\frac{\sqrt{1 + 4x}}{(2 + x)^3}$ in ascending powers of x . [4]**

**(c) Determine the range of values of x for which the
expansion in part (b) is valid, giving a reason for
your answer. [2]**

- 9 (a) Express $3 \cos 2x + 4 \sin 2x$ in the form $R \cos(2x - \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$. [3]
- (b) In this question you must show detailed reasoning.

Solve the equation $\frac{3 \cos 2x + 4 \sin 2x + 6}{3 \cos 2x + 4 \sin 2x - 1} = 4$
for $0^\circ < x < 360^\circ$. [7]

- 10 The graph of $y = e^x$ can be transformed to the graph of $y = e^{2x-1}$ by a stretch parallel to the x -axis FOLLOWED BY a translation.

- (a) (i) State the scale factor of the stretch. [1]
- (ii) Give full details of the translation. [2]

Alternatively the graph of $y = e^x$ can be transformed to the graph of $y = e^{2x-1}$ by a stretch parallel to the x -axis and a stretch parallel to the y -axis.

- (b) State the scale factor of the stretch parallel to the y -axis. [1]

The point P lies on the curve $y = e^{2x-1}$ and has x -coordinate of $\frac{1}{2}$.

- (c) Show that the tangent to the curve $y = e^{2x-1}$ at P has equation $y = 2x$. [4]
- (d) Find the EXACT area enclosed by the curve $y = e^{2x-1}$, the tangent to the curve at P and the y -axis. [4]

- 11 A student is attempting to prove that $\sqrt{5}$ is irrational. The first three lines of their proof are shown below.

Assume that $\sqrt{5}$ is rational, so it can be written as $\sqrt{5} = \frac{a}{b}$. LINE 1

Squaring both sides gives $5 = \frac{a^2}{b^2}$. Hence $a^2 = 5b^2$. LINE 2

Hence a must be a multiple of 5, so $a = 5k$, for some integer k . LINE 3

- (a) State any conditions required on LINE 1. [2]
- (b) Explain why LINE 2 means that a must be a multiple of 5. [2]
- (c) Complete the proof to show that $\sqrt{5}$ is irrational. [3]

- 12 (a) Express $\frac{2 + 4x}{x(1 + x)(1 - x)}$ in partial fractions. [4]

The gradient of a curve is given by

$\frac{dy}{dx} = \frac{2 + 4x}{x(1 + x)(1 - x)\tan y}$ and the curve passes through the point $(\frac{1}{2}, \frac{1}{4}\pi)$.

- (b) Show that the equation of the curve can be written in the form $\cos y = f(x)$, where $f(x)$ is fully simplified. [7]

END OF QUESTION PAPER



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.