

Modified Enlarged 24pt
OXFORD CAMBRIDGE AND RSA EXAMINATIONS

Thursday 19 June 2025 – Afternoon

A Level Mathematics B (MEI)

**H640/03 Pure Mathematics
and Comprehension**

Insert

**Time allowed: 2 hours
plus your additional time allowance**

INSTRUCTIONS

**Do NOT send this Insert for marking. Keep
it in the centre or recycle it.**

INFORMATION

**This Insert contains the article for
Section B.**




The trisectrix of Maclaurin

To bisect an angle means to divide it into two equal angles. It is possible to bisect an angle using a ruler and compasses. The construction is illustrated in FIG. C1.

5

FIG. C1



To trisect an angle means to divide into three equal angles. It is impossible to trisect an angle using a ruler and compasses alone but mathematicians found other methods.

10

In 1742, Colin Maclaurin first studied a curve which can be used to trisect an angle. The curve is called the Trisectrix of Maclaurin.

15

The equation of the curve in cartesian form is $y^2 = \frac{x^2(3a-x)}{a+x}$, where a is a constant. The curve is shown in FIG. C2 on page 4. Only values of $a > 0$ are considered in this article.


20

It can be shown that everywhere on the curve $\frac{3a-x}{a+x} \geq 0$. It follows that $3a-x \geq 0$ and $a+x > 0$.

The curve has the trisection property illustrated in FIG. C2.

25

FIG. C2



The point C has coordinates $(2a, 0)$ and Q is the point where the curve crosses the positive x -axis. The point P is a general point (x, y) on the

30

loop of the curve. The origin of the coordinate system is denoted by O.

The dashed line is an asymptote to the curve.

If the line CP makes an angle 3θ with the positive x -axis then the line OP makes an angle θ with the positive x -axis. Angles are measured anticlockwise from the positive x -axis. 35

To use the curve to trisect an angle, place the angle in position QCP. One arm of the angle (CQ) lies along the x -axis. The other arm of the angle meets the curve at the point P. Angle QOP is one third of angle QCP. 40 45

Deriving the cartesian equation of the trisectrix

Using the formula for $\tan(A + B)$ in terms of $\tan A$ and $\tan B$, it can

be shown that $\tan 3\theta = \frac{t(3 - t^2)}{1 - 3t^2}$, 50

where $t = \tan \theta$.

From this and FIG. C2 it follows

that $\frac{x - 2a}{y} = \frac{1 - 3t^2}{t(3 - t^2)}$.

Hence $\frac{1}{t} - \frac{2a}{y} = \frac{1 - 3t^2}{t(3 - t^2)}$.

It then follows that

55

$$\frac{2a}{y} = \frac{1}{t} - \frac{1 - 3t^2}{t(3 - t^2)} = \frac{2 + 2t^2}{t(3 - t^2)}$$

so $\frac{a}{y} = \frac{1 + t^2}{t(3 - t^2)}$ and this gives

the parametric equation

for y , $y = \frac{at(3 - t^2)}{1 + t^2}$.

The equation for x follows from

60

$\tan \theta = \frac{y}{x}$. Together with the

parametric equation for y , this leads

to $x = \frac{a(3 - t^2)}{1 + t^2}$.

The parameter t can be eliminated from the parametric equations to derive the cartesian equation

65

$$y^2 = \frac{x^2(3a - x)}{a + x}, \text{ as stated in line 17.}$$

BLANK PAGE



Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.